

Mizoroki-Heck cross-couplings of 2-acetyl-5-bromobenzofuran and aryl halides under microwave irradiation

Mohamed R. Shaaban,^{a,b} Ahmed F. Darweesh,^a Kamal M. Dawood^{a,*}
and Ahmad M. Farag^a

^a*Department of Chemistry, Faculty of Science, Cairo University, Giza 12613, Egypt*

^b*Department of Chemistry, Faculty of Applied Science, Umm Al-Qura University,
Makkah Almukarramah , Saudi Arabia*

E-mail: dr_dawood@yahoo.com

DOI: <http://dx.doi.org/10.3998/ark.5550190.0011.a18>

Abstract

Mizoroki-Heck cross-coupling reactions of 2-acetyl-5-bromobenzofuran as well as activated and deactivated aryl bromides with various olefins were investigated under both thermal as well as microwave irradiating conditions in open air using water solvent.

Keywords: Palladium, catalysis, microwave, cross coupling reactions, benzofurans, aryl halides

Introduction

Palladium-catalyzed cross-coupling reactions leading to the formation of carbon-carbon bonds are among the most powerful transformations in organic synthesis.^{1,2} Mizoroki-Heck cross-coupling reaction is a powerful and widely used method to couple aryl halides with terminal olefins to form cinnamates and stilbenes which are of industrial importance.³ The possibility of carrying out Mizoroki-Heck cross-coupling reactions in water and under open air could be of great interest for the development of industrial processes.^{4,5} One of the challenges that are facing organic chemists now is focused on the design of new methodologies able to make the already known chemical transformations simpler, faster, cheaper, and in general, more efficient processes. In particular, increasing attention has been paid to the 'green chemistry' of these processes; this concept being understood as a set of principles⁶ that reduce or eliminate the use or generation of hazardous substances.^{7,8} Furthermore, microwave irradiation methodology assists in achieving rapid incorporation of organic synthesis into broad research diversities.^{9,10}

Intense efforts have been made directed towards the synthesis of benzofuran derivatives due to their biological and pharmacological properties,^{11,12} and to their natural occurrence.^{13,14} In continuation of our recent research work directed to the use of Pd (II)-complexes as precatalysts

in C-C cross couplings in aqueous media,^{15,16} we study here the catalytic reactivity of the Pd(II)-complex¹⁷ **3** in the Mizoroki-Heck cross coupling reactions of 2-acetyl-5-bromobenzofuran as well as aryl and pyridyl halides with a variety of terminal olefins under thermal as well as microwave irradiating conditions. To the best of our knowledge, these are the first Heck vinylation reactions of 2-acetyl-5-bromobenzofuran.

Results and Discussion

Heck cross-coupling reactions of aryl halides


At the beginning, the effect of concentration of the Pd-complex **3** on the Heck cross-coupling between styrene **2** and *p*-bromoacetophenone **1** under thermal heating condition was evaluated. The coupling reaction was conducted in DMF (3 mL) at 130 °C for 90 min using different concentrations of the precatalyst **3** and in all cases the molar ratios of *p*-bromoacetophenone **1**/styrene **2**/ TBAB/ Et₃N: 1/ 1.5/ 0.6/ 3. Therefore, use of 1 mol% of the complex **3** resulted in full conversion into 4-acetylstilbene **4** in 96% isolated yield. The reaction was repeated with different catalytic concentrations; 0.75, 0.5, 0.25, 0.125, 0.05, 0.025, 0.0125 and 0.0 mol% of complex **3** with respect to *p*-bromoacetophenone **1**. Full conversions were obtained at concentrations; 0.75, 0.5, 0.25 and 0.125 mol% of the complex **3** with turnover numbers (TON) 133, 200, 400 and 800 respectively. The reaction was again repeated using 0.05, 0.025 and 0.0125 mol% of the complex **3** giving 4-acetylstilbene **4** in 93, 67 and 13 % GC-conversions with TON 1860, 2680 and 1000, respectively. These results reveal the high activity of the catalytic system that is used in this Heck cross-coupling reaction. Finally, the starting substrates were completely recovered unchanged in the absence of the precatalyst **3**.

Pd-complexes are reported to serve as “dormant species”⁹ that are not involved in the real catalytic cycle but are a source of a catalytically active species of unknown nature. Therefore, the Pd(II)-complex **3** may serve here as a reservoir that is not involved in the real catalytic cycle but is a source of release a considerable amount of colloidal Pd(0) which can show catalytic activity at low concentrations. De Vries and co-workers noted for Heck reactions that low concentrations of Pd(0) in solution will aggregate and deactivate much more slowly than high concentrations.¹⁸ In order to prevent deactivation (precipitation of palladium black) of these active species at higher reaction temperatures, it is necessary to add activating agents such as TBAB to the reaction mixture which has a beneficial effect in Heck reactions due to the ability of TBAB to stabilize Pd(0) nanoparticles against further aggregation and deactivation.¹⁹ The study of TONs in palladium-mediated C-C bond formation processes is a very important area of research and any catalyst displaying >10³ turnover number is defined as high turnover catalyst (HTC).²⁰ Therefore, Pd-complex **3** can be considered as high turn over catalyst.

Next, the catalytic activity of the Pd(II) complex **3** towards Heck cross-coupling reaction of *p*-bromoacetophenone with styrene in different solvents (water, DMF or toluene) and different bases (KOH, K₂CO₃ or Et₃N) using 0.25 mol% under thermal heating was examined. Among the

solvents tested, water as an eco-friendly solvent provided full conversion (85% isolated yield), in the presence of KOH as base and TBAB as additive, after 90 min of thermal heating at 100 °C (entry 1, Table 1). Under this condition, using K₂CO₃ and Et₃N as bases gave 90 and 45% GC-conversions, respectively (entries 3 and 4, Table 1). When the reaction was performed in DMF/Et₃N/TBAB at reflux temperature for 90 min, full conversion with 92% isolated yield was obtained. No coupling product was detected when the last catalytic system was applied at r.t. for 24h. Using DMF as solvent with KOH as base in the presence of TBAB resulted in 89% GC-conversion into 4-acetylstilbene **4** (entry 5, Table 1). Toluene was not a suitable solvent for this reaction regardless the used base (entries 9-11, Table 1). The TLC and ¹H NMR spectrum of the crude reaction product **4**, were more clear and less contaminated with by-products when DMF was the reaction solvent (run 6, Table 1) than when water was used as a solvent (run 1, Table 1), in addition to the higher isolated yield (92%) after run 6 comparing to that obtained after run 1 (85%). Therefore, DMF was the proper solvent for this reaction. The catalytic system (DMF/TBAB/Et₃N) was applied for the same cross coupling reaction under microwave irradiation instead of thermal heating and the reaction was completed within 5 min to give **4** in 88% yield.

Table 1. Effect of base and solvent on Heck coupling of *p*-bromoacetophenone with styrene



Entry	Solvent	Base	Yield% ^{a,b}
1	H ₂ O/TBAB	KOH	100 (85)
2	H ₂ O	KOH	85 (80)
3	H ₂ O/TBAB	K ₂ CO ₃	90 (78)
4	H ₂ O/TBAB	Et ₃ N	45
5	DMF/TBAB	KOH	89
6	DMF/TBAB	Et ₃ N	100 (92) ^{c,d}
7	DMF	Et ₃ N	99 (89)
8	DMF/TBAB	K ₂ CO ₃	97
9	Toluene	KOH	0
10	Toluene	Et ₃ N	11
11	Toluene	K ₂ CO ₃	22

^aConditions: *p*-Bromoacetophenone: 1 mmol; styrene: 1.5 mmol; TBAB: 0.6 mmol; base: 3 mmol; solvent: 3 mL, Pd-complex: 0.25 mol% at reflux condition for 90 min. ^bConversions were based on GC-analysis and the values in parentheses refer to the isolated yields. ^cRepeating this condition at r.t. for 24h resulted in no reaction based on GC-analysis. ^dRepeating this condition under microwave heating (250 Watt) at 160 °C for 5 min resulted full conversion with 88% isolated yield after 5 min.

The optimized catalytic condition using DMF/TBAB/Et₃N under microwave irradiation was applied for carrying out Mizoroki-Heck reactions of activated and deactivated aryl bromides **1**, **5** and **6** with various olefins **2a-d** as outlined in Table 2. Thus, microwave-assisted Heck cross-coupling of the aryl bromides **1**, **5** and **6** with styrene **2a** resulted in full conversion into the corresponding stilbenes **4**, **17** and **21** in very good isolated yields (entries 1, 5 and 9, Table 2). Repeating the latter Heck cross-coupling reactions under thermal heating at 130 °C in DMF/TBAB/ Et₃N for 90 min gave 100% GC-conversion of the corresponding stilbenes **4**, **17** and **21** in 92, 97 and 82% isolated yields, respectively.

Further, arylation of acrylonitrile **2b**, ethyl acrylate **2c** and with *n*-butyl acrylate **2d** with aryl bromides **1**, **5** and **6** proceeded smoothly under microwave irradiation at 160 °C to furnish, in all cases, the corresponding cinnamates **14-16**, **18-20** and **22-24**, respectively, in high isolated yields after a reaction time 4~7 min. The high conversion in the cross-coupling of the less active 4-bromoanisole **6** with the olefins **2a-d** (runs 9-12, Table 2) within 7 min of microwave irradiation gives a good support for the high catalytic activity of the Pd complex **3**. It was reported that strong bases such as KOH decompose *n*-butyl acrylate to give *n*-butanol and acrylic acid²¹ and hence was not used in this study.

Interestingly, the cross-coupling arylation process was highly regio- and stereoselective and provided only the thermodynamically more stable *trans*-isomer of stilbenes, cinnamitriles and cinnamates **4**, **14-24**, where GC, GC-MS and ¹H NMR of the crude reaction mixture showed no evidence for other isomers. The stereochemistry of the double bond in the reaction products was easily established. For example, the protons around the double bond were found to resonate within the aromatic region with a coupling constant about 16 Hz attributed to related olefinic hydrogens.

Table 2. Heck cross-coupling of aryl bromides with olefins under microwave irradiation

Entry	Y	X	Product	MW heating ^{a,b}	
				time(min)	yield%
1	CH ₃ CO	Ph	4	5	100 (88) ^c
2	CH ₃ CO	CN	14	5	100 (93)
3	CH ₃ CO	CO ₂ Et	15	5	100 (78)
4	CH ₃ CO	CO ₂ Bu ⁿ	16	5	100 (85)
5	NO ₂	Ph	17	4	100 (75) ^d
6	NO ₂	CN	18	4	100 (86)
7	NO ₂	CO ₂ Et	19	4	100 (95)
8	NO ₂	CO ₂ Bu ⁿ	20	4	100 (88)
9	OCH ₃	Ph	21	7	100 (87) ^e
10	OCH ₃	CN	22	7	98 (88)
11	OCH ₃	CO ₂ Et	23	7	95 (70)
12	OCH ₃	CO ₂ Bu ⁿ	24	7	90 (81)









^aConditions: Aryl bromide: 1 mmol; olefin: 1.5 mmol; TBAB: 0.6 mmol; Et₃N: 3 mmol; DMF: 3 mL, Pd-complex: 0.25 mol% under microwave heating (250 Watt) at 160 °C. ^bConversions were based on GC-analysis and the values in parentheses refer to the isolated yields. ^cRepeating this reaction under thermal heating at reflux for 90 min gave full conversion with 92% isolated yield. ^dThermal heating for 90 min gave 97% isolated yield. ^eThermal heating for 90 min gave 82% isolated yield.

Next, the applicability of complex **3** in Heck cross-coupling reaction of further activated and deactivated aryl and pyridyl bromides **7-13** with styrene **2a** were carried out under thermal heating as well as microwave irradiation conditions was investigated as described in Table 3. In all cases, styrene was arylated smoothly resulting in the formation of the corresponding stilbenes **25-29** and styrylpyridines **30, 31** in good to excellent isolated yields. As can be seen in Table 3, high chemoselectivity was observed in the case of 1-bromo-4-chlorobenzene (entry 2), where the more reactive bromine atom was replaced with styryl group to give 4-chlorostilbene **26** with 100% conversion under microwave irradiation. Similar cross coupling of 1-bromonaphthalene under the same conditions afforded the cross coupled product **28** in excellent isolated yield with full GC-conversion under both thermal as well as microwave irradiation (entry 4, Table 3). In addition, the Pd(II)-complex **3** was found to efficiently catalyze the coupling of 2-amino-5-bromopyridine **12** and 2-amino-6-bromopyridine **13** to give the corresponding 5-styryl- and 6-styrylpyridines **30** and **31**, respectively, in good yields under thermal and microwave heating modes (entries 6 and 7, Table 3). 2-Amino-6-bromopyridine **12** consumed longer reaction time

than 2-amino-5-bromopyridine **13** under both thermal- and MW-heating reflecting the high activity of 5-bromopyridine than 6-bromopyridine. In all cases, the products were established to have the *trans*-configuration of 1,2-disubstituted ethylenes. The other regioisomers; 1,1-disubstituted ethylenes due to α -arylation²² were not detected in the crude reaction products. In addition, a recent publication reported the partial or complete hydrogenation of the double bond when the reaction was carried out at high temperature, however we could not detect any hydrogenated product during this work.²³


Aryl chlorides are cheaper and more easily available than bromides and iodides, but less reactive. Efficient catalyst systems for the Heck reactions of aryl chlorides have been extensively studied.²⁴ Herein, the reactivity of the Pd-complex **3** (0.25 mol%) in the Heck coupling of activated aryl chlorides (4-chloroacetophenone **32** and 4-chloronitrobenzene **33**) with styrene **2a** was studied under thermal heating as well as microwave irradiation conditions as shown in Scheme 1. Under this condition, very good yields of 4-acetylstilbene **4** and 4-nitrostilbene **17** were obtained. It is noteworthy to mention here that the catalytic activity of the Pd-complex **3** is more efficient in the Heck coupling of aryl chlorides if compared with similar reactions using our previously reported analogous Pd(II)-complexes.¹⁵

Table 3. Heck cross-coupling of aryl and pyridyl bromides with styrene

					
Entry	Ar-Br	Product	Thermal ^a time (h) yield% ^b	Microwave ^a time(min) yield% ^c	
1		25	4 (80)	8	100 (83)
2		26	4 (83)	8	100 (87)
3		27	4 (78)	6	98 (90)
4		28	3 (96)	6	100 (97)
5		29	3 (85)	6	100 (81)
6		30	14 (70)	20	72 (66)
7		31	8 (79)	12	86 (82)

^aConditions: Aryl bromide: 1 mmol; styrene: 1.5 mmol; TBAB: 0.6 mmol; Et₃N: 3 mmol; DMF: 3 mL, Pd-complex: 0.25 mol% under microwave heating (250 Watt) at 160 °C and thermal

heating at 130 °C. ^bAll values refer to the isolated yields. ^cConversions were based on GC-analysis and the values in parentheses refer to the isolated yields.




Scheme 1. Heck cross-coupling of activated aryl chlorides with styrene. Reaction conditions: ^aThermal heating at 130 °C for 6h. ^bMicrowave heating at 160 °C and 250 Watt for 10 min. All values refer to isolated yields.

Heck cross-coupling of 2-acetyl-5-bromobenzofuran

In the next section, we examined the catalytic activity of the Pd(II) complex **3** in the Heck cross-coupling reaction of 2-acetyl-5-bromobenzofuran **34** with styrene **2a** under various reaction conditions as outlined in Table 4. The reaction was carried out in different solvent such as water, dimethylformamide and toluene using Et₃N and KOH as examples of organic and inorganic bases and TBAB as additive was used in order to achieve good yields. In this aspect, water/KOH/TBAB and water/Et₃N/TBAB catalytic systems provided good results 91 and 95% GC-conversions, respectively, into (*E*)- 2-acetyl-5-(β-styryl)benzofuran **35**, (entries 1 and 2, Table 4). The use of DMF/Et₃N/TBAB as catalytic system (entry 5) was more efficient resulting in full conversion with excellent isolated yield (99%) of the product **35** after 4 h of thermal heating at 130 °C using 0.25 mol% of Pd-complex **3**. This condition (DMF/Et₃N/TBAB) was repeated under microwave heating at 160 °C (250 Watt) for 20 min to obtain 100% GC-conversion (92% isolated yield) of **35**, and at room temperature (about 30 °C) for 24 h with stirring to give no reaction at all based on GC-analysis with full recovering of the bromide **34**. On the other hand, when the same reaction was carried out using DMF/KOH/TBAB catalytic system for 4 h of thermal heating it gave lower GC-conversion (85%) than in the case of using DMF/Et₃N/TBAB (entry 4, Table 4). Toluene was found to be not appropriate for cross coupling of the bromide **34** with styrene **2a** in either KOH or Et₃N as bases to give either nil or 4% GC-conversions, respectively (entries 7 and 8, Table 4).

The presence of TBAB is essential for carrying out Heck cross-coupling reactions¹⁹ as shown in Table 4 (entries 1, 2, 4 and 5) and its absence resulted in sharp decrease of the coupling product regardless the used solvent (water or DMF) in the reaction between 2-acetyl-5-bromobenzofuran **34** and styrene under thermal heating (entries 3 and 6, Table 4).

Table 4. Base and solvent effect on Heck coupling of 2-acetyl-5-bromobenzofuran **34** with styrene **2a**


Entry	Solvent	Base	thermal heating ^{a,b} time (h)	yield%
1	H ₂ O/TBAB	KOH	4	91 (89)
2	H ₂ O/TBAB	Et ₃ N	4	95
3	H ₂ O	KOH	4	63
4	DMF/TBAB	KOH	4	85
5	DMF/TBAB	Et ₃ N	4	100 (99) ^{c,d}
6	DMF	Et ₃ N	12	30
7	Toluene	KOH	4	0
8	Toluene	Et ₃ N	4	4

^aConditions: Bromide: 1 mmol; styrene: 1.5 mmol; TBAB: 0.6 mmol; base: 3 mmol; solvent: 3 mL, Pd-complex **3**: 0.25 mol% at reflux condition for 4h. ^bConversions were based on GC-analysis and the values in parenthesis refer to the isolated yields. ^cRepeating this condition at r.t. for 24h resulted in no reaction. ^dRepeating this condition under MW at 160 °C (250 Watt) resulted in 92% yield with full conversion after 20 min.

Based on the foregoing results, vinylation of 2-acetyl-5-bromobenzofuran **34** with acrylic systems such as acrylonitrile **2b**, ethyl acrylate **2c**, *n*-butyl acrylate **2d**, *tert*-butyl acrylate **2e** and methyl acrylate **2f** under both microwave irradiation and thermal heating is illustrated in Table 5. For the synthesis of *E*-3-(2-acetylbenzofuran-5-yl)acrylonitrile **36** from the cross-coupling of **34** with **2b**, the reaction was conducted in DMF/Et₃N/TBAB as catalytic system using 0.25 mol% of the Pd-complex **3**. Full conversion was obtained after 3 hours of thermal heating (isolated yield 88%) and after 20 min of microwave irradiation (isolated yields 88%) of the benzofuranylacrylonitrile **36** (entry 1, Table 5). Under similar catalytic cross-coupling condition, 2-acetyl-5-bromobenzofuran **34** was vinyated with further acrylates **2c-f** and the Heck coupling reaction was completed within 3-4 hours under thermal heating and after 20 min of microwave irradiation with high isolated yields (>80%) of the corresponding 5-(benzofuranyl)acrylates **37-40** (Table 5, entries 2, 3, 4, 6). The role of TBAB as an additive was crucial for the effective vinylation in the synthesis of the cross-coupled products **36-40**, however in the absence of TBAB a dramatic effect appeared as no conversion at all after 16 h of thermal heating or 30 min of microwave irradiation during the coupling of *tert*-butyl acrylate **2e** with 2-acetyl-5-bromobenzofuran **34** (Table 5, entry 5). The same effect for absence of TBAB was encountered in the coupling of methyl acrylate **2f** with **34** giving only 17% conversion after 16 h of thermal

heating (entry 7, Table 5) reflecting the essential role of TBAB in such cross coupling reactions. The structures of the newly synthesized cross-coupled products **36-40** were established on the basis of their elemental and spectral analyses as described in the experimental section.

Table 5. Heck cross-coupling of 2-acetyl-5-bromobenzofuran **34** with acrylates **2b-f**

Reaction scheme: R-CH=CH_2 (**2b-f**) + **34** $\xrightarrow[\text{0.25 mol\% Cat. 3}]{\text{DMF/Et}_3\text{N/TBAB}}$ **36-40**

Entry	R	Product No.	Thermal heating ^{a,b}		MW heating ^{a,b}	
			time (h)	yield%	time(min)	yield%
1	CN	36	3	100 (88)	20	100 (88)
2	CO ₂ Et	37	3	100 (91)	20	100 (81)
3	CO ₂ Bu ⁿ	38	3	100 (80)	20	100 (91)
4	CO ₂ Bu ^t	39	4	100 (86)	20	100 (95)
5	CO ₂ Bu ^t	39	16 ^c	0	30	0
7	CO ₂ Me	40	4	100 (93)	n.d.	n.d.
8	CO ₂ Me	40	16 ^c	17	n.d.	n.d.

^aConditions: Bromide: 1 mmol; olefin: 1.5 mmol; TBAB: 0.6 mmol; Et₃N: 3 mmol; DMF: 3 mL, Pd-complex: 0.25 mol%, microwave heating (250 Watt) at 160 °C and thermal heating at 130 °C.

^bConversions were based on GC-analysis and the values in parentheses refer to the isolated yields. ^cWithout TBAB. n.d.= not done

In conclusion, the Pd(II)-complex **3** was found to be efficient and highly active precatalyst for Mizoroki-Heck cross-coupling reactions of activated and deactivated aryl halides with styrene and acrylates under thermal heating as well as microwave irradiation conditions. Complex **3** was also highly active precatalyst when applied to the first vinylation reactions of 2-acetyl-5-bromobenzofuran. In all studied cases, 0.25 mol% of the Pd-complex **3** were sufficient for full conversion within short reaction times.

Experimental Section

General. Melting points were determined in open glass capillaries with a Gallenkamp apparatus. The infrared spectra were recorded in potassium bromide disks on a Pye Unicam SP 3-300 and Shimaduz FTIR 8101 PC infrared spectrophotometer. NMR spectra were recorded with a Varian

Mercury VXR-300 NMR spectrometer at 300 MHz (^1H NMR) and at 75 MHz (^{13}C NMR) using CDCl_3 as solvent and internal standard (δ 7.27 and 77.36 ppm, for ^1H NMR and ^{13}C NMR, respectively). Capillary GC analyses were performed with a Shimadzu GC-14A or GC-14B, a Shimadzu C-R6A integrator, and an HP 5 column (25 m length, 0.25 mm i.d., 0.25 μm film) or recorded with an Agilent GC 6890N. Mass spectra (EI) were obtained at 70 eV with a type Shimadzu GCMQP 1000 EX spectrometer. Analytical thin-layer chromatography was performed using pre-coated silica gel 60778 plates (Fluka), and the spots were visualized with UV light at 254 nm. Fluka silica gel 60741 (70-230 mesh) was used for flash column chromatography. Microwave experiments were carried out using a CEM Discover LabmateTM microwave apparatus (300 W with ChemDriverTM Software).

Effect of concentration of Pd-complex 3 on Heck coupling of 4-bromoacetophenone with styrene in DMF under thermal heating

A mixture of 4-bromoacetophenone **1** (199 mg, 1 mmol) and styrene **2a** (156 mg, 1.5 mmol), TBAB (194 mg, 0.6 mmol), palladium complex **3** (4.59 mg, 1 mol%), TEA (303 mg, 3 mmol) and DMF (3 mL) was stirred at 130 °C under open air for 1.5 h to give 4-acetylstilbene **4**. The same experiment was repeated using of palladium complex **3** in different mol% *e.g.* 0.75, 0.5, 0.25, 0.125, 0.05, 0.025, 0.0125 and 0.0 mol% of **3** with respect to 4-bromoacetophenone **1**. The molar ratios of the reaction components were in all cases as follows; 4-bromoacetophenone, styrene, TBAB, TEA, DMF: 1 / 1.5 / 0.6 / 3 / 3 mL DMF.

Effect of base and solvent on Heck coupling of 4-bromoacetophenone with styrene under thermal heating

A mixture of 4-bromoacetophenone **1** (199 mg, 1 mmol) and styrene **2a** (156 mg, 1.5 mmol), TBAB (194 mg, 0.6 mmol), Pd-complex **3** (1.14 mg, 0.25 mol%), and potassium hydroxide (168 mg, 3 mmol) in water (3 mL) was stirred at 130 °C under open air for 1.5 h. After the reaction was almost complete (monitored by GC), to give 4-acetylstilbene **4**. The same experiment was repeated using different solvent and base. The molar ratios of the reaction components were in all cases as follows; 4-bromoacetophenone, styrene, TBAB, base, and solvent: 1 / 1.5 / 0.6 / 3 / 3 mL. The yield % versus different solvents and bases is outlined in Table 1.

Heck vinylation of aryl bromides in DMF under microwave heating

A mixture of the appropriate aryl bromides **1**, **5** or **6** (1 mmol) and the appropriate olefins **2a-d** (1.5 mmol), TBAB (0.6 mmol), Pd-complex **3** (0.25 mol%), and Et_3N (303 mg, 3 mmol) in DMF (3 mL) was mixed in a process vial. The vial was capped properly, and the mixture was then heated under microwave irradiating conditions at 160 °C and 250 Watt for the appropriate reaction time as listed in Table 2. After the reaction was almost complete (monitored by TLC), the reaction mixture was left to cool to room temperature. The reaction mixture was then extracted three times with EtOAc (3 x 20 mL) and then the organic fractions were combined together, dried over MgSO_4 , filtered and then the solvent was removed under vacuum. The

residue was then subjected to purification via flash column chromatography with petroleum *n*-hexane–EtOAc (10:1) as eluent to give the corresponding pure products **4** and **14-24**.

(E)-4-Acetylstilbene (4). White crystals; mp 134-136 °C (Ref.²⁵ mp 134-136 °C); ¹H NMR (CDCl₃) δ 2.61 (s, 3H, CH₃CO), 7.13 (d, 1H, *J* = 16.2 Hz), 7.24 (d, 1H, *J* = 16.2 Hz), 7.29-7.42 (m, 3H), 7.55 (d, 2H, *J* = 7.5 Hz), 7.59 (d, 2H, *J* = 8.4 Hz), 7.96 (d, 2H, *J* = 8.1 Hz); MS *m/z* (%) 222 (68.7, M⁺), 176 (37.3), 151 (25.4), 115 (43.3), 91 (100), 75 (32.8), 56 (40.3).

(E)-4-Acetylcinnamonitrile (14). Buff crystals; mp 84-86 °C (Ref.²⁶ mp is not mentioned); ¹H NMR (CDCl₃) δ 2.56 (s, 3H, CH₃CO), 5.97 (d, 1H, *J* = 16.8 Hz), 7.38 (d, 1H, *J* = 16.8 Hz), 7.51 (d, 2H, *J* = 8.1 Hz), 7.93 (d, 2H, *J* = 8.1 Hz); MS *m/z* (%) 171 (19, M⁺), 157 (15.5), 115 (100), 93 (29.3), 73 (60.3), 55 (96.6).

(E) Ethyl 4-acetylcinnamate (15). White crystals; mp 40-42 °C (Ref.²⁷ mp 42-44 °C); ¹H NMR (CDCl₃) δ 1.28 (t, 3H, *J* = 7.2 Hz), 2.54 (s, 3H, CH₃CO), 4.21 (q, 2H, *J* = 7.2 Hz), 6.45 (d, 1H, *J* = 15.9 Hz), 7.52 (d, 2H, *J* = 8.4 Hz), 7.62 (d, 1H, *J* = 15.9 Hz), 7.89 (d, 2H, *J* = 8.4 Hz); MS *m/z* (%) 218 (23.5, M⁺), 203 (68.6), 175 (34), 156 (26.1), 115 (100), 91 (34.6), 73 (66), 55 (99.3).

(E) *n*-Butyl 4-acetylcinnamate (16). A colorless oil;²⁸ ¹H NMR (CDCl₃) δ 0.88 (t, 3H, *J* = 7.2 Hz), 1.31-1.39 (m, 2H), 1.58-1.63 (m, 2H), 2.50 (s, 3H, CH₃CO), 4.13 (t, 2H, *J* = 6.6 Hz), 6.43 (d, 1H, *J* = 16.2 Hz), 7.50 (d, 2H, *J* = 8.1 Hz), 7.58 (d, 1H, *J* = 16.2 Hz), 7.86 (d, 2H, *J* = 8.1 Hz); MS *m/z* (%) 246 (20.2, M⁺), 190 (60), 156 (24.3), 115 (100), 93 (32.5), 73 (70.1), 55 (97.8).

(E)-4-Nitrostilbene (17). Yellow crystals; mp 160-162 °C (Ref.²⁹ mp 161-162 °C); ¹H NMR (CDCl₃) δ 7.15 (d, 1H, *J* = 16.2 Hz), 7.28 (d, 1H, *J* = 16.2 Hz), 7.34-7.44 (m, 3H), 7.56 (d, 2H, *J* = 8.1 Hz), 7.64 (d, 2H, *J* = 8.7 Hz), 8.22 (d, 2H, *J* = 8.7 Hz); MS *m/z* (%) 225 (100, M⁺), 178 (9.1), 153 (13.6), 119 (15.9), 65 (36.4), 50 (79.5).

(E)-4-Nitrocinnamonitrile (18). Brown crystals; mp 198-200 °C (Ref.³⁰ mp 198-199 °C); ¹H NMR (CDCl₃) δ 6.06 (d, 1H, *J* = 16.5 Hz), 7.47 (d, 1H, *J* = 16.5 Hz), 7.63 (d, 2H, *J* = 9 Hz), 8.27 (d, 2H, *J* = 9 Hz); MS *m/z* (%) 174 (35.3, M⁺), 130 (54.4), 102 (69.8), 76 (20.5), 55 (100).

(E) Ethyl 4-nitrocinnamate (19). Pale yellow crystals; mp 132-134 °C (Ref.²⁷ mp 133-136 °C); ¹H NMR (CDCl₃) δ 1.36 (t, 3H, *J* = 7.2 Hz), 4.30 (q, 2H, *J* = 7.2 Hz), 6.56 (d, 1H, *J* = 16.2 Hz), 7.67 (d, 2H, *J* = 8.1 Hz), 7.71 (d, 1H, *J* = 16.2 Hz), 8.25 (d, 2H, *J* = 8.1 Hz); MS *m/z* (%) 221 (33.3, M⁺), 176 (100), 130 (63.2), 102 (64.9), 77 (22.8), 63 (54.4), 50 (19.3).

(E) *n*-Butyl 4-nitrocinnamate (20). Pale yellow crystals; mp 62-64 °C (Ref.³¹ mp 63-65 °C); ¹H NMR (CDCl₃) δ 0.97 (t, 3H, *J* = 6.9 Hz), 1.39-1.48 (m, 2H), 1.66-1.73 (m, 2H), 4.24 (t, 2H, *J* = 6.6 Hz), 6.56 (d, 1H, *J* = 15.9 Hz), 7.67 (d, 2H, *J* = 9 Hz), 7.71 (d, 1H, *J* = 15.6 Hz), 8.24 (d, 2H, *J* = 8.7 Hz); MS *m/z* (%) 249 (5, M⁺), 194 (30.2), 176 (60.6), 130 (24.7), 102 (75.2), 90 (32), 76 (53.6), 56 (100).

(E)-4-Methoxystilbene (21). Buff powder; mp 134-136 °C (Ref.³² mp 134-136 °C); ¹H NMR (CDCl₃) δ 3.85 (s, 3H, OCH₃), 6.93 (d, 2H, *J* = 8.7 Hz), 7.01 (d, 1H, *J* = 16.5 Hz), 7.11 (d, 1H, *J* = 16.5 Hz), 7.24-7.30 (m, 1H), 7.35-7.40 (m, 2H), 7.48 (d, 2H, *J* = 8.7 Hz), 7.52 (d, 2H, *J* = 7.8 Hz); MS *m/z* (%) 210 (100, M⁺), 195 (68.3), 179 (61), 88 (43.9), 74 (64.3), 64 (46.3).

(E)-4-Methoxycinnamonitrile (22). Pale yellow powder; mp 60-62 °C (Ref.³³ mp 62-63 °C); ¹H NMR (CDCl₃) δ 3.83 (s, 3H, OCH₃), 5.69 (d, 1H, *J* = 16.5 Hz), 6.90 (d, 2H, *J* = 8.1 Hz), 7.31 (d,

1H, $J = 16.5$ Hz), 7.38 (d, 1H, $J = 8.1$ Hz); MS m/z (%) 159 (100, M^+), 145 (63.3), 108 (49.2), 73 (59.4), 55(39).

(E) Ethyl 4-methoxycinnamate (23). Yellowish white crystals; mp 48-50 °C (Ref.³¹ mp 48-50 °C); ¹H NMR (CDCl₃) δ 1.32 (t, 3H, $J = 7.2$ Hz), 3.80 (s, 3H, OCH₃), 4.24 (q, 2H, $J = 7.2$ Hz), 6.29 (d, 1H, $J = 15.9$ Hz), 6.88 (d, 2H, $J = 7.8$ Hz), 7.45 (d, 2H, $J = 8.1$ Hz), 7.63 (d, 1H, $J = 15.9$ Hz); MS m/z (%) 206 (75.1, M^+), 183 (43.7), 149 (39.3), 108 (36.9), 71 (34.2), 57 (100).

(E) *n*-Butyl 4-methoxycinnamate (24). Colorless oil;²⁸ ¹H NMR (CDCl₃) δ 0.95 (t, 3H, $J = 7.2$ Hz), 1.39-1.46 (m, 2H), 1.65-1.70 (m, 2H), 3.80 (s, 3H, OCH₃), 4.19 (t, 2H, $J = 6.9$ Hz), 6.30 (d, 1H, $J = 15.9$ Hz), 6.87 (d, 2H, $J = 6.9$ Hz), 7.44 (d, 2H, $J = 6.9$ Hz), 7.62 (d, 1H, $J = 15.9$ Hz); MS m/z (%) 234 (27.6, M^+), 178 (56.1), 156 (49.5), 108 (43.4), 75 (36.4), 57 (100).

Heck coupling of styrene with aryl halides in DMF under thermal heating

A mixture of the appropriate aryl halide **7-13**, **32** and **33** (1 mmol) and styrene **2a** (1.5 mmol), TBAB (0.6 mmol), Pd-complex **3** (0.25 mol%), and Et₃N (303 mg, 3 mmol) in DMF (3 mL) was stirred at 130 °C under open air for the appropriate reaction time as listed in Table 3. After the reaction was almost complete (monitored by TLC), the reaction mixture was left to cool to room temperature. The reaction mixture was then extracted three times with EtOAc (3 x 20 mL) and then the organic fractions were combined together, dried over MgSO₄, filtered and then the solvent was removed under vacuum. The residue was then subjected to purification via flash column chromatography with petroleum *n*-hexane–EtOAc (10:1) as eluent to give the corresponding pure products **25-31**, **4** and **17**, respectively.

Heck coupling of styrene with aryl halides in DMF under microwave heating

A mixture of the appropriate aryl halide **7-13**, **32** and **33** (1 mmol) and styrene **2a** (1.5 mmol), TBAB (0.6 mmol), Pd(II)-complex **3** (0.25 mol%), and TEA (303 mg, 3 mmol) in DMF (3 mL) was mixed in a process vial. The vial was capped properly, and the mixture was then heated under microwave irradiating conditions at 160 °C and 250 Watt for the appropriate reaction time as listed in Table 3. After the reaction was almost complete (monitored by TLC), the reaction mixture was left to cool to room temperature. The reaction mixture was then extracted three times with EtOAc (3 x 20 mL) and then the organic fractions were combined together, dried over MgSO₄, filtered and then the solvent was removed under vacuum. The residue was then subjected to purification via flash column chromatography with *n*-hexane–EtOAc (10:1) as eluent to give the corresponding pure products **25-31**, **4** and **17**, respectively.

(E)-4-Ethoxystilbene (25). Pale yellow crystals; mp 122-124 °C (Ref.³⁴ mp 124-126 °C); ¹H NMR (CDCl₃) δ 1.42 (t, 3H, $J = 7.2$ Hz), 4.01 (q, 2H, $J = 7.2$ Hz), 6.78 (d, 2H, $J = 9$ Hz), 6.86-6.92 (m, 1H), 6.98 (d, 1H, $J = 16.5$ Hz), 7.09 (d, 1H, $J = 16.5$ Hz), 7.38 (d, 2H, $J = 9$ Hz), 7.45-7.48 (m, 2H), 7.51 (d, 2H, $J = 6.9$ Hz); MS m/z (%) 224 (17.3, M^+), 167 (25.8), 149 (48.9), 129 (30.8), 104 (32.0), 91 (34.4), 77 (67.1), 57 (100).

(E)-4-Chlorostilbene (26). Buff crystals; mp 128-130 °C (Ref.³² mp 128-129 °C); ¹H NMR (CDCl₃) δ 7.04 (d, 1H, $J = 16.8$ Hz), 7.11 (d, 1H, $J = 16.8$ Hz), 7.28-7.35 (m, 3H), 7.38 (d, 2H, J

= 7.2 Hz), 7.45 (d, 2H, J = 8.4 Hz), 7.51 (d, 2H, J = 7.5 Hz); MS m/z (%) 214 (100) (M^+), 178 (58.3), 115 (44.4), 91 (66.7), 75 (55.6), 50 (61.1).

(E)-3-Acetylstilbene (27). Colorless crystals; mp 78-80 °C (Ref.³³ mp 79-80 °C); ^1H NMR (CDCl_3) δ 2.65 (s, 3H, COCH_3), 7.14 (d, 1H, J = 16.5 Hz), 7.21 (d, 1H, J = 16.5 Hz), 7.29-7.34 (m, 1H), 7.35-7.41 (m, 2H), 7.44-7.49 (m, 1H), 7.54 (d, 2H, J = 7.5 Hz), 7.71 (d, 1H, J = 7.2 Hz), 7.84 (d, 1H, J = 7.8 Hz), 8.11 (s, 1H); MS m/z (%) 222 (58.3, M^+), 179 (45.8), 51 (100).

(E)-1-Styrylnaphthalene (28). Colorless oil;³⁵ ^1H NMR (CDCl_3) δ 7.30 (d, 1H, J = 16.2 Hz), 7.41-7.70 (m, 6H), 7.74 (d, 2H, J = 7.8 Hz), 7.88 (d, 1H, J = 7.2 Hz), 7.93 (d, 1H, J = 8.4 Hz), 7.99-8.07 (m, 2H), 8.38 (d, 1H, J = 8.1 Hz); MS m/z (%) 230 (100, M^+), 229 (94.3), 215 (24.3), 152 (43.5), 114 (29), 101 (22.4), 45 (32.8).

(E)-4-Benzoylstilbene (29). White crystals; mp 148-150 °C (Ref.³⁶ mp 148-149 °C); IR (KBr) ν 1644 cm^{-1} (C=O); ^1H NMR (CDCl_3) δ 7.17 (d, 1H, J = 16.2 Hz), 7.26 (d, 1H, J = 16.2 Hz), 7.31-7.53 (m, 6H), 7.56 (d, 2H, J = 7.5 Hz), 7.62 (d, 2H, J = 8.7 Hz), 7.80-7.85 (m, 4H); MS m/z (%) 284 (65.1, M^+), 207 (55.2), 178 (60.4), 105 (51), 77 (100).

(E)-2-Amino-6-styrylpyridine (30). Orange crystals; mp 106-108 °C (Ref.³⁷ mp 106-108 °C); ^1H NMR (CDCl_3) δ 4.89 (s, 2H, NH_2), 6.36 (d, 1H, pyridine-3H, J = 8.1 Hz), 6.75 (d, 1H, pyridine-5H, J = 7.2 Hz), 7.03-7.11 (m, 1H, pyridine-4H), 7.17 (d, 1H, J = 16.2), 7.22-7.30 (m, 3H), 7.32 (d, 2H, J = 8.1 Hz), 7.38 (d, 1H, J = 16.2); MS m/z (%) 196 (12, M^+), 181 (23.2), 167 (27.2), 149 (52.7), 129 (36), 104 (31.5), 91 (55.4), 77 (75.8), 57 (100).

(E)-2-Amino-5-styrylpyridine (31). Yellow crystals; mp 200-202 °C (Ref.³⁸ mp 199-200 °C); ^1H NMR (CDCl_3) δ 4.51 (s, 2H, NH_2), 6.53 (d, 1H, pyridine-3H, J = 8.4 Hz), 6.92 (d, 1H, J = 16.2 Hz), 7.07 (d, 1H, J = 16.2 Hz), 7.22-7.38 (m, 3H), 7.48 (d, 2H, J = 7.8 Hz), 7.68 (d, 1H, pyridine-4H, J = 8.7 Hz), 8.18 (s, 1H, pyridine-6H); MS m/z (%) 196 (50, M^+), 178 (52.8), 139 (55.6), 115 (66.7), 93 (44.4), 69 (100), 57 (63.9).

Effect of base and solvent on Heck coupling of 2-acetyl-5-bromobenzofuran (34) with styrene under thermal heating

A mixture of 2-acetyl-5-bromobenzofuran **34** (239 mg, 1 mmol) and styrene **2a** (156 mg, 1.5 mmol), TBAB (194 mg, 0.6 mmol), Pd-complex **3** (1.14 mg, 0.25 mol%), and potassium hydroxide (168 mg, 3 mmol) in water (3 mL) was stirred at 130 °C under open air for 4 h. After the reaction was almost complete (monitored by GC), to give 2-acetyl-5-(β -styryl)benzofuran **35**. The same experiment was repeated using different solvent and base. The molar ratio of the reaction components were in all cases as follows; 2-acetyl-5-bromobenzofuran **34**, styrene **2a**, TBAB, base, solvent: 1 / 1.5 / 0.6 / 3 / 3 mL. The yield % versus different solvents and bases is outlined in Table 4.

Heck coupling of 2-acetyl-5-bromobenzofuran (34) with acrylates in DMF under thermal heating

A mixture of 2-acetyl-5-bromobenzofuran **34** (239 mg, 1 mmol) and appropriate acrylate **2b-f** (1.5 mmol), TBAB (0.6 mmol), complex **3** (0.25 mol%), and Et_3N (303 mg, 3 mmol) in DMF (3 mL) was stirred at 130 °C under open air for the appropriate reaction time as listed in Table 5.

After the reaction was almost complete (monitored by TLC), the reaction mixture was left to cool to room temperature. The reaction mixture was then extracted three times with EtOAc (3 x 20 mL) and then the organic fractions were combined together, dried over MgSO₄, filtered and then the solvent was removed under vacuum. The residue was then subjected to purification via flash column chromatography with petroleum *n*-hexane–EtOAc (7:1) as eluent to give the corresponding pure products of 5-(benzofuranyl)acrylates **36-40**.

Heck coupling of 2-acetyl-5-bromobenzofuran (34) with acrylates in DMF under microwave heating

A mixture of 2-acetyl-5-bromobenzofuran **34** (239 mg, 1 mmol) and appropriate olefin **2b-f** (1.5 mmol), TBAB (0.6 mmol), complex **3** (0.25 mol%), and Et₃N (303 mg, 3 mmol) in DMF (3 mL) was mixed in a process vial. The vial was capped properly, and the mixture was then heated under microwave irradiating conditions at 160 °C and 250 Watt for 20 min. After the reaction was almost complete (monitored by TLC), the reaction mixture was left to cool to room temperature. The reaction mixture was then extracted three times with EtOAc (3 x 20 mL) and then the organic fractions were combined together, dried over MgSO₄, filtered and then the solvent was removed under vacuum. The residue was then subjected to purification via flash column chromatography with petroleum *n*-hexane–EtOAc (7:1) as eluent to give the corresponding pure products 5-(benzofuranyl)acrylates **36-40**.

(E)-2-Acetyl-5-(β -styryl)benzofuran (35). Yellow powder; mp. 187-189 °C; *R_f* 0.54 (hexane : AcOEt 7:1); IR (KBr) ν 3119 (w), 3024 (w), 1676 (s), 1297 (m), 1074 (s) cm⁻¹; ¹H NMR (CDCl₃) δ 2.63 (s, 3H, CH₃CO), 7.11 (d, 1H, *J* = 16.5 Hz), 7.21 (d, 1H, *J* = 16.5 Hz), 7.25-7.31 (m, 1H), 7.36-7.41 (m, 2H), 7.50-7.58 (m, 4H), 7.69 (d, 1H, *J* = 9 Hz), 7.80 (s, 1H); ¹³C NMR (DEPT) (CDCl₃) δ 26.4 (1CH₃), 112.6, 112.9, 120.9, 126.4, 126.9, 127.7, 128.0, 128.7, 128.8 (9CH), 127.5, 133.7, 137.1, 153.2, 155.3, 188.5 (6C); MS *m/z* (%) 262 (100, M⁺), 247 (14.5), 191 (34.6), 95 (10.6), 63 (10.0), 51 (11.0). Anal. Calcd for C₁₈H₁₄O₂: C, 82.42; H, 5.38. Found: C, 82.33; H, 5.27%.

(E)-3-(2-Acetylbenzofuran-5-yl)acrylonitrile (36). Yellow powder; mp. 122-124 °C; *R_f* 0.57 (hexane : AcOEt 7:1); IR (KBr) ν 3113 (w), 3046 (w), 2205 (m), 1684 (m), 1552 (s), 1276 (s) cm⁻¹; ¹H NMR (CDCl₃) δ 2.63 (s, 3H, CH₃CO), 5.91 (d, 1H, *J* = 16.5 Hz), 7.51 (d, 1H, *J* = 16.5 Hz), 7.51 (s, 1H), 7.59-7.62 (m, 2H), 7.80 (s, 1H); ¹³C NMR (DEPT) (CDCl₃) δ 26.5 (1CH₃), 96.2, 112.5, 113.4, 123.1, 126.7, 150.0 (6CH), 118.0, 127.7, 129.8, 153.8, 156.7, 188.4 (6C); MS *m/z* (%) 211 (72.9, M⁺), 196 (100), 184 (22.0), 140 (44.1), 93 (16.9), 75 (33.9), 58 (64.4). Anal. Calcd for C₁₃H₉NO₂: C, 73.92; H, 4.29; N, 6.63. Found: C, 74.01; H, 4.16; N, 6.55%.

(E) Ethyl 3-(2-acetylbenzofuran-5-yl)acrylate (37). Yellow powder; mp. 108-110 °C; *R_f* 0.66 (hexane : AcOEt 7:1); IR (KBr) ν 3102 (w), 2927 (w), 1710 (m), 1671 (s), 1635 (m), 1283 (m), 1171 (s) cm⁻¹; ¹H NMR (CDCl₃) δ 1.36 (t, 3H, CO₂CH₂CH₃, *J* = 7.2 Hz), 2.63 (s, 3H, CH₃CO), 4.29 (q, 2H, CO₂CH₂CH₃, *J* = 7.2 Hz), 6.46 (d, 1H, *J* = 16.2 Hz), 7.51 (s, 1H), 7.59 (d, 1H, *J* = 9 Hz), 7.68 (d, 1H, *J* = 8.7 Hz), 7.78 (d, 1H, *J* = 15.9 Hz), 7.85 (s, 1H); ¹³C NMR (DEPT) (CDCl₃) δ 14.3, 26.5 (2CH₃), 60.6 (1CH₂), 112.8, 113.0, 118.2, 123.5, 127.7, 144.0 (6CH), 127.6, 130.7,

153.5, 156.4, 166.8, 188.4 (6C); MS m/z (%) 258 (100, M^+), 213 (92.9), 186 (36.8), 171 (56.1), 115 (30.8), 63 (39.5). Anal. Calcd for $C_{15}H_{14}O_4$: C, 69.76; H, 5.46. Found: C, 69.65; H, 5.32%.

(E) Butyl 3-(2-acetylbenzofuran-5-yl)acrylate (38). Yellow powder; mp. 76-78 °C; R_f 0.57 (hexane : AcOEt 7:1); IR (KBr) ν 3097 (w), 2956 (w), 2872 (w), 1709 (m), 1680 (s), 1634 (m), 1284 (m), 1170 (m) cm^{-1} ; 1H NMR ($CDCl_3$) δ 0.98 (t, 3H, $J = 7.5$ Hz), 1.42-1.49 (m, 2H), 1.66-1.73 (m, 2H), 2.63 (s, 3H, CH_3CO), 4.23 (t, 2H, $J = 6.6$ Hz), 6.46 (d, 1H, $J = 15.9$ Hz), 7.51 (s, 1H), 7.59 (d, 1H, $J = 8.7$ Hz), 7.68 (d, 1H, $J = 8.7$ Hz), 7.78 (d, 1H, $J = 15.9$ Hz), 7.85 (s, 1H); ^{13}C NMR (DEPT) ($CDCl_3$) δ 13.7, 26.5 (2 CH_3), 19.1, 30.7, 64.5 (3 CH_2), 112.8, 113.0, 118.2, 123.5, 127.7, 143.9 (6CH), 127.6, 130.7, 153.5, 156.4, 166.9, 188.4 (6C); MS m/z (%) 286 (25.4, M^+), 230 (100), 213 (65.5), 171 (36.6), 115 (23.7), 56 (23.5). Anal. Calcd for $C_{17}H_{18}O_4$: C, 71.31; H, 6.34. Found: C, 71.26; H, 6.21%.

(E) tert-Butyl 3-(2-acetylbenzofuran-5-yl)acrylate (39). White powder; mp. 150-152 °C; R_f 0.58 (hexane : AcOEt 6:1); IR (KBr) ν 3119 (w), 3004 (w), 2927 (w), 1706 (m), 1670 (m), 1631 (m), 1285 (m), 1080 (m) cm^{-1} ; 1H NMR ($CDCl_3$) δ 1.47 (s, 9H, $C(CH_3)_3$), 2.55 (s, 3H, CH_3CO), 6.31 (d, 1H, $J = 15.9$ Hz), 7.43 (s, 1H), 7.50 (d, 1H, $J = 8.7$ Hz), 7.59 (d, 1H, $J = 9$ Hz), 7.61 (d, 1H, $J = 15.6$ Hz), 7.76 (s, 1H); ^{13}C NMR (DEPT) ($CDCl_3$) δ 26.5, 28.2 (2 CH_3), 112.8, 112.9, 120.1, 123.3, 127.7, 142.9 (6CH), 80.6, 127.6, 131.0, 153.4, 156.3, 166.1, 188.4 (7C); MS m/z (%) 286 (8.3, M^+), 230 (100), 215 (91.4), 171 (6.5), 114 (8.3), 57 (13.7). Anal. Calcd for $C_{17}H_{18}O_4$: C, 71.31; H, 6.34. Found: C, 71.32; H, 6.18%.

(E) Methyl 3-(2-acetylbenzofuran-5-yl)acrylate (40). Yellow powder; mp. 146-148 °C; R_f 0.55 (hexane : AcOEt 6:1); IR (KBr) ν 3108 (w), 2952 (w), 1706 (m), 1674 (s), 1631 (m), 1283 (s), 1143 (s) cm^{-1} ; 1H NMR ($CDCl_3$) δ 2.55 (s, 3H, CH_3CO), 3.75 (s, 3H, CO_2CH_3), 6.38 (d, 1H, $J = 15.9$ Hz), 7.44 (s, 1H), 7.51 (d, 1H, $J = 8.7$ Hz), 7.61 (d, 1H, $J = 8.7$ Hz), 7.72 (d, 1H, $J = 16.2$ Hz), 7.77 (s, 1H); ^{13}C NMR (DEPT) ($CDCl_3$) δ 26.5, 51.7 (2 CH_3), 112.7, 113.0, 117.7, 123.6, 127.7, 144.3 (6CH), 127.6, 130.7, 153.5, 156.4, 167.3, 188.4 (6C); MS m/z (%) 244 (100, M^+), 229 (46.0), 213 (62.1), 171 (27.5), 115 (17.9), 63 (10.0). Anal. Calcd for $C_{14}H_{12}O_4$: C, 68.85; H, 4.95. Found: C, 68.95; H, 4.90%.

Acknowledgements

K.M.D and his student A.F.D are very thankful to Prof. Peter Metz (TU-Dresden) for giving them the chance to measure GC-analysis at his laboratory during the A.v.H. fellowships; July-Sept. 2008. Also, K.M.D. is deeply indebted to A.v.H. Foundation for the gift of a CEM Discover LabmateTM microwave apparatus.

References

1. (a) Malleron, J. L.; Fiaud, J. C.; Legros, J. Y. *Handbook of Palladium-Catalysed Organic Reactions*; Academic: San Diego, CA, 1997. (d) Dupont, J.; Pfeffer, M.; Spencer, J. *Eur. J. Inorg. Chem.* **2001**, 1917. (b) M. Beller, A. Zapf: In *Handbook of Organopalladium Chemistry for Organic Synthesis*, Vol. 1 Negishi, E.-I., Ed., Wiley-Interscience: New York, 2002; pp 1209-1222. (c) Beller, M.; Bolm, C. *Transition Metals for Organic Synthesis*, 2nd Edn., Wiley-VCH: Weinheim, 2004. (d) de Meijere, A.; Diederich, F. *Metal-Catalyzed Cross-Coupling Reactions*, 2nd Edn., Wiley-VCH: Weinheim, 2004.
2. (a) Beller, M.; Zapf, A. *Top. Catal.* **2002**, *19*, 101. (b) Kantchev, E. A. B.; O'Brien, C. J.; Organ, M. G. *Angew. Chem. Int. Ed.* **2007**, *46*, 2768.
3. For Reviews see: (a) Heck, R. F. *Acc. Chem. Res.* **1979**, *12*, 146. (b) Cabri, W.; Candiani, I. *Acc. Chem. Res.* **1995**, *28*, 2. (c) Negishi, E.; Coperet, C.; Ma, S.; Liou, S.; Liu, F. *Chem. Rev.* **1996**, *96*, 365. (d) Crisp, G. T. *Chem. Soc. Rev.* **1998**, *27*, 427. (e) Beletskaya, J. P.; Cheprakov, A. V. *Chem. Rev.* **2000**, *100*, 3009. (f) Biffis, A.; Zecca, M.; Basato, M. *J. Mol. Catal. A* **2001**, *173*, 249. (g) Whitcombe, N. J.; Hii, K. K.; Gibson, S. E. *Tetrahedron* **2001**, *57*, 7449. (h) Dounay, A. B.; Overman, L. E. *Chem. Rev.* **2003**, *103*, 2945. (i) Alonso, F.; Beletskaya, I. P.; Yus, M. *Tetrahedron* **2005**, *61*, 11771.
4. (a) De Meijere, A.; Meyer, F. *Angew. Chem., Int. Ed. Engl.* **1994**, *33*, 2379. (b) Beletskaya, I. P.; Cheprakov, A. V. *Chem. Rev.* **2000**, *100*, 3009. (c) Blaser, H.-U. *Chem. Commun.* **2003**, 293. (d) Blaser, H.-U.; Indolese, A.; Naud, F.; Nettekoven, U.; Schnyder, A. *Adv. Synth. Catal.* **2004**, *346*, 1583.
5. (a) Withcombe, N.; Hii-Mimi, K. K.; Gibson, S. *Tetrahedron* **2001**, *57*, 7449. (b) Botella, L.; Nájera, C. *Tetrahedron* **2004**, *60*, 5563. (c) Botella, L.; Nájera, C. *J. Org. Chem.* **2005**, *70*, 4360.
6. Anastas, P. T.; Warner, J. C. *Green Chemistry: Theory and Practice*; Oxford University Press: New York, 1998; p 30.
7. (a) Anastas, P. T.; Williamson, T. C. *Green Chemistry: Designing Chemistry for the Environment*; ACS: Washington, DC, 1996. (b) Anastas, P. T.; Heine, P. T.; Williamson, T. C. Eds. *Green Chemical Synthesis and Processes*; ACS: Washington, DC, 2000. (c) Lancaster, M. *Green Chemistry: An Introductory Text*; RSC: Cambridge, 2002.
8. (a) Larhed, M.; Moberg, C.; Hallberg, A. *Acc. Chem. Res.* **2002**, *35*, 717. (b) Desai, B.; Kappe, C. O. In: *Immobilized Catalysts* Kirschning, A., Ed. *Top. Curr. Chem.* **2004**, *242*, 177. (c) Kappe, C. O. *Angew. Chem. Int. Ed.* **2004**, *43*, 6250. (d) Hoz, A.; Ortiz, A. D.; Moreno, A. *Chem. Soc. Rev.* **2005**, *34*, 164.
9. (a) Mingos, D. M. P.; Baghurst, D. R. *Chem. Soc. Rev.* **1991**, *20*, 1. (b) Strauss, C. R.; Trainor, R. W. *Aust. J. Chem.* **1995**, *48*, 1665. (c) Gabriel, C.; Gabriel, S.; Grant, E. H.; Halstead, B. S.; Mingos, D. M. P. *Chem. Soc. Rev.* **1998**, *27*, 213.

10. (a) Elander, N.; Jones, J. R.; Lu, S. Y.; Stone-Elander, S. *Chem. Soc. Rev.* **2000**, 29, 239. (b) Lidström, P.; Tierney, J.; Wathey, B.; Westman, J. *Tetrahedron* **2001**, 57, 9225. (c) Larhed, M.; Hallberg, A. *Drug Discovery Today* **2001**, 6, 406.
11. (a) Smith, R. A.; Chen, J.; Mader, M. M.; Muegge, I.; Moehler, U.; Katti, S.; Marrero, D.; Stirtan, W. G.; Weaver, D. R.; Xiao, H.; Carley, W. *Bioorg. Med. Chem. Lett.* **2002**, 12, 2875. (b) Hayakawa, I.; Shioya, R.; Agatsuma, T.; Furukawa, H.; Sugano, Y. *Bioorg. Med. Chem. Lett.* **2004**, 14, 3411. (c) Cowart, M.; Faghieh, R.; Curtis, M. P.; Gfesser, G. A.; Bennani, Y. L.; Black, L. A.; Pan, L.; Marsh, K. C.; Sullivan, J. P.; Esbenshade, T. A.; Fox, G. B.; Hancock, A. A. *J. Med. Chem.* **2005**, 48, 38. (d) Hall, A.; Billinton, A.; Brown, S. H.; Clayton, N. M.; Chowdhury, A.; Giblin, G. M. P.; Goldsmith, P.; Hayhow, T. G.; Hurst, D. N.; Kilford, I. R.; Naylor, A.; Passingham, B.; Winyard, L. *Bioorg. Med. Chem. Lett.* **2008**, 18, 3392.
12. (a) Dawood, K. M.; Abdel-Gawad, H.; Ragab, E. A.; Ellithy, M.; Mohamed, H. A. *Bioorg. Med. Chem.* **2006**, 14, 3672; (b) Dawood, K. M.; Abdel-Gawad, H.; Ellithy, M.; Mohamed, H. A.; Hegazi, B. *Arch. Pharm. Chem. Life Sci.* **2006**, 339, 133; (c) Abdel-Aziz, H. A.; Mekawey, A. A. I.; Dawood, K. M. *Eur J. Med. Chem.* **2009**, 44, 3637.
13. (a) Halabalaki, M.; Aligiannis, N.; Papoutsis, Z.; Mitakou, S.; Moutsatsou, P.; Sekeris, C.; Skaltsounis, A. *J. Nat. Prod.* **2000**, 63, 1672; (b) Apers, S.; Paper, D.; Bürgermeister, J.; Baronikova, S.; Dyck, S. V.; Lemie' re, G.; Vlietinck, A.; Pieters, L. *J. Nat. Prod.* **2002**, 65, 718; (c) Stevenson, P. C.; Simmonds, M. S. J.; Yule, M. A.; Veitch, N. C.; Kite, G. C.; Irwin, D.; Legg, M. *Phytochemistry* **2003**, 63, 41.
14. (a) Akgul, Y. Y.; Anil, H. *Phytochemistry* **2003**, 63, 939; (b) Ali, Z.; Tanaka, T.; Iliya, I.; Iinuma, M.; Furusawa, M.; Ito, T.; Nakaya, K.-i.; Murata, J.; Darnaedi, D. *J. Nat. Prod.* **2003**, 66, 558; (c) Kokialakis, N.; Magiatis, P.; Mitaku, S.; Pratsinis, H.; Tilleguin, F. *Planta Med.* **2003**, 69, 566; (d) Aslam, S. N.; Stevenson, P. C.; Phythian, S. J.; Veitch, N. C.; Hall, D. R. *Tetrahedron* **2006**, 62, 4214.
15. (a) Dawood, K. M.; Kirschning, A. *Tetrahedron* **2005**, 61, 12121; (b) Dawood, K. M. *Tetrahedron* **2007**, 63, 9642; (c) Dawood, K. M.; Fayed, M. S.; Elkhalea, M. M. *Arkivoc* **2009** (xiii), 324; (d) Dawood, K. M.; Eldeftar, M. M. *Synthesis*, **2010**, 1030.
16. (a) Solodenko, W.; Brochwitz, C.; Wartchow, R.; Hashem, Md. A.; Vaultier, M.; Dawood, K. M.; Kirschning, A. *Mol. Div.* **2005**, 9, 333; (b) Dawood, K. M.; Solodenko, W.; Kirschning, A. *Arkivoc* **2007** (v), 104.
17. Darweesh, A. F.; Shaaban, M. R.; Farag, A. F.; Metz, P.; Dawood, K. M. *Synthesis*, **2010**, accepted for publication.
18. De Vries, A. H. M.; Mulders, J. M. C. A.; Mommers, J. H. M.; Henderickx, H. J. W.; de Vries, J. G. *Org. Lett.*, **2003**, 5, 3285.
19. (a) Jeffery, T. *Adv. Met. Org. Chem.* 1996, 153; (b) Zapf, A.; Beller, M. *Chem. Eur. J.* **2001**, 7, 2908; (c) Reetz, M. T.; Westermann, E. *Angew. Chem. Int. Ed.* **2000**, 39, 165.
20. Farina, V. *Adv. Synth. Catal.* **2004**, 346, 1553.
21. Sud, A.; Deshpande, R. M.; Chaudhari, R. V. *J. Mol. Catal. A: Chem.* **2007**, 270, 144.

22. (a) Gruendemann, S.; Albrecht, M.; Loch, J. A.; Faller, J. W.; Crabtree, R. H. *Organometallics* **2001**, *20*, 5485; (b) Loch, J. A.; Albrecht, M.; Peris, E.; Mata, J.; Faller, J. W.; Crabtree, R. H. *Organometallics* **2002**, *21*, 700.
23. Malik, I.; Hussain, M.; Ali, A.; Toguem, S.-M. T.; Basha, F. Z.; Fischer, C.; Langer, P. *Tetrahedron* **2010**, *66*, 1637.
24. (a) Schnyder, A.; Aemmer, J.; Indolese, A. F.; Pittelkow, U.; Studer, M. *Adv. Synth. Catal.* **2002**, *344*, 495. (b) Diez-Barra, E.; Guerra, J.; Hornillos, V.; Merino, S.; Tejeda, J. *Organometallics* **2003**, *22*, 4610. (c) Consorti, C. S.; Zanini, M. L.; Leal, S.; Ebeling, G.; Dupont, J. *Org. Lett.* **2003**, *5*, 983; (d) Consorti, C. S.; Ebeling, G.; Flores, F. R.; Rominger, F.; Dupont, J. *Adv. Synth. Catal.* **2004**, *346*, 617. (e) Yang, D.; Chen, Y.-C.; Zhu, N.-Y. *Org. Lett.* **2004**, *6*, 1577. (f) Prockl, S. S.; Kleist, W.; Gruber, M. A.; Kohler, K. *Angew. Chem., Int. Ed.* **2004**, *43*, 1881. (g) Frey, G. D.; Schutz, J.; Herdtweck, E.; Herrmann, W. A. *Organometallics* **2005**, *24*, 4416; (h) Prockl, S.; Kleist, W.; Kohler, K. *Tetrahedron* **2005**, *61*, 9855.
25. Kantam, M. L. *Tetrahedron* **2007**, *63*, 8002.
26. Pathak, D. D.; Maheswaran, H.; Prasanth, K. L.; Kantam, M. L. *Synlett* **2007**, *5*, 757.
27. Mu, B. *Tetrahedron* **2007**, *63*, 11475.
28. Wang, R.; Twamley, B.; Shreeve, J. M. *J. Org. Chem.* **2006**, *71*, 426.
29. Suzuki, T. *J. Am. Chem. Soc.* **2005**, *127*, 11720.
30. Cai, M. Z. *Chin. Chem. Lett.* **2005**, *16*, 449.
31. Li, J. -H. *Eur. J. Org. Chem.* **2006**, *22*, 5099.
32. Aggarwal, V. K. *J. Am. Chem. Soc.* **2003**, *125*, 6034.
33. Gokel, G. W. *Tetrahedron Lett.* **1976**, *39*, 3495.
34. Heynekamp, J. J. *J. Med. Chem.* **2006**, *49*, 7182.
35. Alacid, E. *J. Org. Chem.* **2008**, *73*, 2315.
36. Listvan, V. N. *Zh. Obshchei Khim.* **1980**, *50*, 1523.
37. Honma, Y.; Hanamoto, K.; Hashiyama, T.; Sekine, Y.; Takeda, M.; Ono, Y.; Tsuzuraharat, K. *J. Med. Chem.* **1984**, *27*, 125.
38. Bumagin, N. A.; Bykov, V. V.; Sukhomlinova, L. I.; Tolstaya, T. P.; Beletskaya, I. P. *J. Organomet. Chem.* **1995**, *486*, 259.