Synthesis and charge-transfer complexation studies on bis(aminomethyl) *m*-terphenyl based bis-oxycyclophanes with intra-annular amide functionality

Perumal Rajakumar^{a,*} and Ramar Padmanabhan^b

^aDepartment of Organic Chemistry, University of Madras, Guindy Campus, Chennai – 600 025, India ^bResearch and Development Centre, Orchid Chemicals and Pharmaceuticals Ltd., Sozhanganallur, Chennai – 600 119, India E-mail: <u>perumalrajakumar@gmail.com</u>

DOI: http://dx.doi.org/10.3998/ark.5550190.0013.614

Abstract

Bis(aminomethyl) *m*-terphenyl based bis-oxy cyclophanes with amide group as intra-annular functionality were synthesized and characterized from spectral and analytical data. All the cyclophanes form 1:1 charge-transfer (CT) complex with TCNQ.

Keywords: Cyclophane amides, bis-oxy amides, charge transfer complex, TCNQ

Introduction

Synthesis of architecturally novel supramolecules in the context of designing simple models for studying supramolecular interactions stimulates the synthetic chemist to modify the molecular structures. Introduction of amide, ester functionality¹⁻⁴ at the annular ring of cyclophanes would make them as models of protein-metal binding sites in biological systems.⁵⁻⁷ Manganese-oxo complexes of porphyrin ligands^{8,9} are reactive intermediates in O-atom transfer process. Cyclophanes containing both ether and amide functionality at the intra-annular ring system show selectivity in the complexation of metal ions.¹⁰ The biphenyl¹¹ based cyclic amides have been reported for anion complexation.¹² Supramolecular amides were used as molecular receptors¹³ and in molecular recognition^{14,15} of biologically interacting substrates including anti-HIV active macrocyclic amides.¹⁶ Synthesis of amide based supramolecular systems has been reported in the literature.¹⁷⁻²⁰ The most important aspect of supramolecular chemistry is the host-guest complexation. Interactions between electron donors and complementary electron acceptor groups in cyclophanes can form intramolecular charge-transfer (CT) complexes and can exhibit self complementary properties in addition to π - π interaction.²¹⁻²³ Hence it is of interest to synthesize

bis-oxy cyclophane diamides **1-5** and study their CT complexation properties with 7,7,8,8-tetracyanoquinodimethane (TCNQ), tetracyanoethylene (TCNE) and paraquat (PQT).



Figure 1. Structures of cyclophane amides 1-5.

Results and Discussion

Seven different bis-oxy cyclophane amides 1-5 shown in Figure 1 were synthesized from a novel bis(aminomethyl) *m*-terphenyl 22. Reaction of 1.0 equiv. of bis(bromomethyl) *m*-terphenyl 21²⁴ with 2.2 equiv. of hexamine²⁵ in chloroform at reflux resulted in the formation of hexammonium salt. Hydrolysis of hexammonium salt with hydrochloric acid in EtOH-H₂O mixture at reflux afforded diamine 22 in about 90% yield (Scheme 1). The structure of diamine 22 was confirmed from the spectral data. The ¹H NMR spectrum of diamine 22 displayed the *N*-methylene protons as a singlet at δ 3.93. The rest of the aromatic protons appeared in the region δ 7.26-7.79.

Reaction of 1.0 equiv. of *o*, *m*, *p*-xylylene dibromide, *m*-nitro xylylene dibromide and 2,6bis(bromomethyl) pyridine with 2.3 equiv. of ethyl *m*-hydroxybenzoate in the presence of 3.1 equiv. potassium carbonate in DMF at room temperature resulted in the formation of the corresponding bis-oxy esters **6**, **6a**, **6b**, **7**, and **8** respectively in about 85-95% yields after purification by column chromatography. Similarly reaction of 1.0 equiv. of resorcinol and



Figure 2. Structures of dicarboxylic acid ester 6-10, dicarboxylic acid 11-15 and dicarboxylic acid chloride 16-20.

2,8-naphthalenediol with 2.2 equiv. of ethyl bromoacetate in the presence of 3.1 equiv. potassium carbonate in acetonitrile in the presence of catalytic amount of KI at reflux for 8 h resulted in the formation of bis-oxy esters 9, and 10 respectively in about 80-95% yields after purification by column chromatography. The structure of bis-oxy esters 6, 6a, 6b, 7, 8, 9 and 10 was confirmed from spectral and analytical data. Hydrolysis of bis-oxy esters with potassium hydroxide in aqueous ethanol at reflux followed by acidification with 6M HCl afforded the corresponding bis-oxy acids 11, 11a, 11b, 12, 13, 14 and 15 in about 90-95% yields. The structure of bis-oxy acids 11, 11a, 11b, 12, 13, 14 and 15 was confirmed from spectral and

analytical data. The corresponding bis-oxy acid chlorides **16**, **16a**, **16b**, **17**, **18**, **19** and **20** were prepared by reacting the corresponding bis-oxy acids with thionyl chloride in the presence of TEA in DCM (Figure 2).



Scheme 1. Reagents and conditions: (i) Hexamine, CHCl₃, rt, 12 h; (ii) Conc.HCl, EtOH-H₂O, reflux, 3 h; (iii) NaOH-H₂O, rt, 22 (90%); (iv) TEA, DCM, rt, 24 h, 1 (35%), 1a (40%), 1b (52%), 2 (42%), 3 (43%), 4 (30%) and 5 (40%).

In order to test the synthetic utility of diamine **22** for the synthesis of bis-oxy cyclophane diamide, 1.0 equiv of diamine **22** was coupled with 1.1 equiv of diacid chloride **16**, **16a**, **16b**, **17**, **18**, **19** and **20** in the presence of triethylamine in dry DCM at room temperature under high dilution conditions. The reaction afforded the bis-oxy cyclophane diamides **1**, **1a**, **1b**, **2**, **3**, **4** and **5** in 35, 40, 52, 42, 43, 30 and 40% yields respectively, after purification by column chromatography (Scheme 1). The ¹H NMR spectrum of cyclophane amide **1a** displayed a doublet for the *N*-methylene protons at δ 4.51, a singlet for *O*-methylene protons at δ 5.13 and NH protons as a triplet at δ 9.03. The rest of the aromatic protons appeared between δ 7.22 and 7.73. In the ¹³C NMR spectrum of cyclophane amide **1a**, the *N*-methylene carbon appeared at δ 42.2, *O*-methylene carbons at δ 69.6 and carbonyl carbons at δ 166.7. The FT-IR spectrum showed the carbonyl stretching frequency at 1639 cm⁻¹ for the cyclophane amide **1a**. Similarly

the structure of the cyclophane amides 1, 1b, 2, 3, 4 and 5 has been confirmed from the spectral and analytical data.

Determination of association constant (Ka) by Benesi-Hildebrand method

The presence of ether and amide functionality in cyclophanes 1-5 can make them good molecular receptors for electron deficient guest molecules like TCNQ, TCNE and PQT. Cyclophane amides 1, 1a, 1b, 2, 3, 4 and 5 exhibited the formation of charge transfer complexes with TCNQ. However, complexation studies of compounds 1, 1a, 1b, 2, 3, 4 and 5 with TCNE and PQT were

Table 1. λ_{max} for cyclophane amides and TCNQ complex of cyclophane amides 1, 1a, 1b, 2, 3, 4 and 5

Conc. of guest,	Absorbance,	[Y]/A(M)	$1/[X] (M^{-1})$
[X] (M)	А		
4.9 x 10 ⁻⁶	0.032	0.000591	204081
9.8 x 10 ⁻⁶	0.051	0.000371	102040
14.7 x 10 ⁻⁶	0.064	0.000295	68027
19.6 x 10 ⁻⁶	0.073	0.000259	51020
24.5 x 10 ⁻⁶	0.080	0.000236	40816
29.4 x 10 ⁻⁶	0.088	0.000215	34013

not successful. However, cyclophanes **1**, **1a**, **1b**, **2**, **3**, **4** and **5** show UV-Vis absorption maxima at 256.0, 256.0, 256.5, 257.0 and 256.0 nm respectively and the acceptor TCNQ shows an absorption maximum at 393.0 nm. Cyclophanes **1**, **1a**, **1b**, **2**, **3**, **4** and **5** form a charge-transfer

Cyclophane	Ka (mol ⁻¹ dm ³)	$\epsilon [M^{-1}cm^{-1}]$	r
amide			
1	$6.52 \ge 10^4$	$4.36 \ge 10^3$	0.9999
1 a	6.65 x 10 ⁴	6.87 x 10 ³	0.9998
1b	$8.75 \ge 10^4$	$4.04 \text{ x } 10^3$	0.9994
2	1.49 x 10 ⁵	4.77 x 10 ³	0.9999
3	$8.06 \ge 10^4$	5.96 x 10 ³	0.9993
4	6.34 x 10 ⁴	6.45×10^3	0.9998
5	1.03×10^5	$4.46 \ge 10^3$	0.9993

Table 2. Complexation of TCNQ with cyclophane amides 1, 1a, 1b, 2, 3, 4 and 5

complex with TCNQ as evidenced by the appearance of absorption maxima at 743.0, 743.0, 742.0, 743.0 and 743.0 nm respectively (Table 1). The plot of (concentration of cyclophane) / absorbance (Y/A) vs 1/concentration of guest (1/X) was linear. Benesi-Hildebrand equation was employed to calculate Ka values. ²⁶ From the slope and the intercept values, K_a (K_a = intercept X

slope⁻¹) and ε (ε = intercept⁻¹) were evaluated. The plot of (concentration of cyclophane) / absorbance (Y/A) vs 1/concentration of guest (1/X) is linear suggesting that the predominate species in solution is a 1:1 complex (Fig. S3). The Ka, ε and r values of the CT complexes formed from **1**, **1a**, **1b**, **2**, **3**, **4** and **5** with TCNQ are shown in Table 2. Benesi-Hildebrand treatment for the CT complex formed between the cyclophane amide **1a** and TCNQ is shown in table 3 and Figure S3. CT spectra for cyclophane amide **1a** and **2** with variable concentration of TCNQ are shown in Figure S1 and S2.

,		
Cyclophane amide	λ_{max} (nm) of	λ_{max} (nm) of TCNQ
	cyclophane amide	complex
1	256.0	743.0
1a	256.0	743.0
1b	256.5	742.0
2	257.0	743.0
3	256.0	743.0
4	255.0	743.5
5	258.0	742.5

Table 3. Benesi-Hildebrand treatment data of the CT complex formed between the cyclophane amide, **1a** and TCNQ

 $\lambda_{max} = 743.5 \text{ nm}; \text{ concentration of cyclophane amide, } \mathbf{1a} = 1.89 \text{ x } 10^{-5} \text{ M}.$ Ka = 6.65 x 10⁴ M⁻¹; $\epsilon = 6.87 \text{ x } 10^3 \text{ [M}^{-1}\text{cm}^{-1}\text{] and } r = 0.9998.$

Conclusions

In summary we have synthesized various cyclophane amides which show strong CT interactions selectively with TCNQ rather than TCNE and PQT. The biological activity and detailed charge transfer complexation studies of other similar cyclophane amides are under investigation.

Experimental Section

General. Melting points were determined by using a Toshniwal melting point apparatus by open capillary tube method and are uncorrected. The UV visible spectra were recorded on Shimadzu 2550 spectrophotometer. The IR spectra were performed on a PerkinElmer series 2000 FT-IR spectrophotometer. ¹H NMR and ¹³ C NMR spectra were recorded on a Bruker Advance 400 NMR spectrophotometer with TMS as internal standard. Mass spectra were determined by ESI-MS using PerkinElmer Sciex, API 3000 spectrophotometer. Elemental analyses were performed on a Perkin-Elmer 240B elemental analyser.

General procedure for the synthesis of bis-oxy cyclophane amides (1-5)

A solution of diamine 22 (1.4 mmol) in dry dichloromethane (200 mL) and a solution of the corresponding diacid chloride 16, 16a, 16b, 17, 18, 19 and 20 (1.54 mmol) in dichloromethane (200 mL) were simultaneously added dropwise to a well stirred solution of triethylamine (3.1 mmol) in dry dichloromethane (400 mL) for 8 h. After the addition was complete the reaction mixture was stirred for another 24 h. The solvent was removed at reduced pressure and the residue obtained was then dissolved in chloroform (300 mL), washed with water (2 x 200 mL) to remove triethylammonium chloride and then dried over anhydrous sodium sulphate. Removal of the chloroform under reduced pressure gave the corresponding cyclophane amide as a crude material, which was purified by column chromatography (SiO₂).

Cyclophane amide (1). Yield 35%, mp 315 °C, IR (KBr, cm⁻¹): 1713, 1639, 1600. ¹H NMR (400 MHz, DMSO-*d*₆) δ 4.43 (d, 4H, *J* = 5.9 Hz), 5.24 (s, 4H), 7.19-7.21 (m, 2H), 7.38-7.46 (m, 13H), 7.49-7.52 (m, 3H), 7.56 (d, 4H, *J* = 8.0 Hz), 7.60 (d, 2H, *J* = 7.0 Hz), 9.06 (t, 2H, *J* = 6 Hz). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 42.7, 67.2, 113.2, 118.7, 119.54, 124.4, 127.2, 128.0, 128.2, 128.9, 129.4, 129.5, 135.1, 136.5, 139.2, 139.6, 141.3, 158.7, 166.4. MS (ES) *m/z*: 631.2 ([M+H]⁺). Anal. Calcd. for C₄₂H₃₄N₂O₄: C, 79.98; H, 5.43; N, 4.44%. Found: C, 80.67; H, 5.47; N, 4.47%.

Cyclophane amide (1a). Yield 40%, mp 260 °C, IR (KBr, cm⁻¹): 1639, 1583, 1539. ¹H NMR (400 MHz, DMSO-*d*₆) δ 4.51 (d, 4H, *J* = 5.9 Hz), 5.13 (s, 4H), 7.21-7.25 (m, 2H), 7.41-7.44 (m, 10H), 7.46-7.58 (m, 5H), 7.63 (d, 1H, *J* = 1.5 Hz), 7.67 (s, 1H), 7.68 (d, 4H, *J* = 8.2 Hz), 7.73 (bs, 1H), 9.03 (t, 2H, *J* = 5.9 Hz). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 42.2, 69.6, 113.8, 116.9, 119.6, 119.9, 124.9, 127.1, 127.7, 128.2, 128.7, 129.1, 129.6, 136.7, 137.0, 138.8, 139.2, 140.9, 158.3, 166.8. MS (ES) *m/z*: 631.2 ([M+H]⁺). Anal. Calcd. for C₄₂H₃₄N₂O₄: C, 79.98; H, 5.43; N, 4.44%. Found: C, 80.75; H, 5.46; N, 4.48%.

Cyclophane amide (1b). Yield 52%, mp 284 °C, IR (KBr, cm⁻¹): 1638, 1583, 1539. ¹H NMR (400 MHz, DMSO-*d*₆) δ 4.52 (d, 4H, *J* = 5.8 Hz), 5.19 (s, 4H), 7.26-7.28 (m, 2H), 7.39-7.45 (m, 10H), 7.48 (s, 4H), 7.53 (d, 1H, *J* = 7.2 Hz), 7.64 (d, 2H, *J* = 7.5 Hz), 7.69 (d, 4H, *J* = 8.1 Hz), 7.80 (bs, 1H), 9.01 (t, 2H, *J* = 5.8 Hz). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 42.0, 68.9, 114.8, 116.2, 119.7, 125.0, 126.5, 127.1, 12.5, 128.1, 129.5, 129.7, 136.5, 138.7, 139.1, 140.8, 158.0, 166.5. MS (ES) *m*/*z*: 631.2 ([M+H]⁺). Anal. Calcd. for C₄₂H₃₄N₂O₄: C, 79.98; H, 5.43; N, 4.44%. Found: C, 80.55; H, 5.48; N, 4.47%.

Cyclophane amide (2). Yield 42%, mp 285 °C, IR (KBr, cm⁻¹): 1641, 1584, 1532. ¹H NMR (400 MHz, DMSO-*d*₆) δ 4.50 (d, 4H, *J* = 5.8 Hz), 5.30 (s, 4H), 7.24-7.27 (m, 2H), 7.40-7.45 (m, 10H), 7.55 (t, 1H, *J* = 7.0 Hz), 7.63-7.67 (m, 7H), 8.06 (s, 1H), 8.39 (s,2H), 9.05 (t, 2H, *J* = 6.0 Hz). MS (ES) *m/z*: 676.4 ([M+H]⁺). Anal. Calcd. for C₄₂H₃₃N₃O₆: C, 74.65; H, 4.92; N, 6.22%. Found: C, 74.86; H, 4.98; N, 6.31%.

Cyclophane amide (3). Yield 43%, mp 247 °C, IR (KBr, cm⁻¹): 1638, 1588, 1536. ¹H NMR (400 MHz, DMSO- d_6) δ 4.48, 4.55 (a pair of distorted doublet, 4H), 5.15 (distorted ABq, 4H), 7.23-7.26 (m, 5H), 7.34-7.54 (m, 11H), 7.65 (d, 2H, J = 7.5 Hz), 7.71 (d, 2H, J = 7.3 Hz), 7.77

(bs, 1H), 7.90-8.05 (m, 2H), 8.94 (t, 2H, J = 8.5 Hz). MS (ES) m/z: 631.7 ([M+H]⁺). Anal. Calcd. for C₄₁H₃₃N₃O₄: C, 77.95; H, 5.27; N, 6.65%. Found: C, 78.16; H, 5.31; N, 6.69%.

Cyclophane amide (4). Yield 30%, mp 300 °C, IR (KBr, cm⁻¹): 1667, 1602, 1535. ¹H NMR (400 MHz, DMSO- d_6) δ 4.37 (distorted doublet, 4H), 4.56 (s, 4H), 6.63 (d, 3H, J = 9.5 Hz), 7.23 (t, 1H, J = 7.6 Hz), 7.30 (d, 4H, J = 7.2 Hz), 7.49 (d, 1H, J = 7.0 Hz) 7.58 (d, 2H, J = 6.9 Hz) 7.65 (d, 4H, J = 7.4 Hz) 7.82 (s, 1H), 8.67 (distorted triplet, 2H). MS (ES) m/z: 496.7 ([M+NH₄]⁺). Anal. Calcd. for C₃₀H₂₆N₂O₄: C, 75.30; H, 5.48; N, 5.85%. Found: C, 75.59; H, 5.53; N, 5.89%.

Cyclophane amide (5). Yield 40%, mp 260 °C, IR (KBr, cm⁻¹): 1742, 1660, 1634, 1531. ¹H NMR (400 MHz, DMSO-*d*₆) δ 4.31 (d, 4H, *J* = 6.1 Hz), 4.55 (s, 4H), 6.49, 6.51 (dd, 2H, *J* = 2.2 Hz), 7.09 (d, 2H, *J* = 2.0 Hz), 7.13 (s, 1H), 7.21 (d, 2H, *J* = 9.0 Hz), 7.43 (ABq, 8H, *J* = 8.0 Hz), 7.55-7.64 (m, 3H), 8.85 (t, 2H, *J* = 6.1 Hz). ¹³C NMR (100 MHz, DMSO-*d*₆) δ : 42.2, 67.5, 109.7, 112.3, 124.1, 124.7, 127.0, 128.5, 128.6, 128.9, 129.5, 135.0, 138.9, 139.5, 141.3, 156.2, 168.2. MS (ES) *m*/*z*: 529.2 ([M+H]⁺). Anal. Calcd. for C₃₄H₂₈N₂O₄: C, 77.25; H, 5.34; N, 5.30%. Found: C, 77.48; H, 5.41; N, 5.38%.

CT complexation studies of cyclophane amides 1, 1a, 1b, 2, 3, 4 and 5 with TCNQ. A solution of TCNQ ($4.9 \times 10^{-6} \text{ M}$) in CH₃CN at various dilutions (1 mL, 2 mL, 3 mL, 4 mL, 5 mL and 6 mL) were prepared and added to the solution of the cyclophane amide 1, 1a, 1b, 2, 3, 4 and 5 ($1.89 \times 10^{-5} \text{ M}$) in a 1:1 mixture of CHCl₃/CH₃CN (3 mL) in a quartz cuvette of path length 1 cm. The UV-Vis spectrum was also obtained for each of the sample separately and the changes in the absorbance of CT bands were recorded.

Supplementary Material

CT spectra of cyclophane amides **1a** and **2** (Figure S1 and S2) and plot between 1/X and Y/A for cyclophane amide **1a** (Figure S3) are available.

Acknowledgements

The authors thank DST, New Delhi for financial assistance, RSIC, CDRI, Lucknow for FAB MS, and DST-FIST for providing NMR facility for the Department. RP thanks Orchid Chemicals & Pharmaceuticals Ltd., for providing the laboratory and analytical facility.

References

1. Krzysztof, E. K.; Jerald, S.B.; Daria, J. Z. Chem. Rev. 1989, 89, 929-972.

- (a) Ulrich, L. Liebigs Ann. Chem. 1987, 11, 949-955. (b) Dirk, L.; Rainer, A. J. B.; Helmar, G. Liebigs Ann. 1994, 1, 199-201.
- 3. (a) Jingsong, Y.; Xiaoqi, Y.; Changlu, L.; Rugang, X. Synth. Commun. 1999, 29, 2447-2455.
 (b) Yukito, M.; Junichi, K.; Hiroaki, T. J. Chem. Soc. Chem. Commun. 1985, 753-755.
- 4. Thomas, R. W.; Yuan, F.; Cynthia, J. B. J. Org. Chem. 1989, 54, 1584-1589.
- 5. Hosseini, M. W.; Lehn, J. M.; Du, S. R.; Gu, K.; Mertes, M. P. J. Org. Chem. 1987, 52, 1662-1666.
- 6. Lehn, J. M. Science (Washington, D.C.). Inclusion Compounds 1985, 227, 849-859.
- 7. Yohannes, P. G.; Mertes, M. P.; Mertes, K. B. J. Am. Chem. Soc. 1985, 107, 8288-8289.
- 8. Groves, J. T.; Stern, M. K. J. Am. Chem. Soc. 1988, 110, 8628-8638.
- 9. Groves, J. T.; Lee, J.; Marla, S. S. J. Am. Chem. Soc. 1997, 119, 6269-6273.
- 10. Lehn, J. M. Supramolecular chemistry: Concepts and Perspectives VCH: Weinheim, Germany, 1995.
- 11. Costero, A. M.; Banuls, M. J.; Aurell, M. J.; Ward, M. D.; Argent, S. *Tetrahedron* **2004**, *60*, 9471-9478.
- 12. (a) Szumna, A.; Jurczak, J. A. *Eur. J. Org. Chem.* **2001**, *21*, 4031-4039. (b) Chmielewski, M.; Jurczak, J. *Tetrahedron Lett.* **2004**, *45*, 6007-6010.
- 13. Christopher, A. H.; Duncan, H. P. Angew. Chem. Int. Ed. Engl. 1992, 31, 792-795.
- 14. Kyu, C. S.; Donna, V. E.; Erkang, F.; Andreq, D. H. J. Am. Chem. Soc. 1991, 113, 7640-7645.
- 15. Claire, M. K.; David, A. L. Angew. Chem. Int. Ed. 2004, 43, 1222-1223.
- 16. Jhaumeer-Laulloo, B. S. Asian J. Chem. 2000, 12, 775-780.
- 17. Subodh, K.; Vandana, B.; Harijit, S. Tetrahedron 1998, 54, 5575-5586.
- 18. Yehia, A.; Ahmed, H. M. Synthesis 1993, 5, 503-508.
- 19. Rajkumar.P.; Abdul Rasheed, A. M.; Balu. P. M.; Murugesan. K. *Bioorg. Med. Chem.* 2006, *14*, 7458-7467.
- 20. Rajkumar.P.; Abdul Rasheed, A. M.; Rabia. A. I.; Chamundeeswari. D. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 6019-6023.
- 21. Bauer, H.; Stier, F.; Petry, C.; Stadler, C.; Staab, H. A. Eur J. Org. Chem. 2001, 3255-3279.
- 22. Sorenson, A. P.; Nielsen. M. B.; Becher, J. Tetrahedron Lett. 2003, 44, 2979-2982.
- 23. (a) Rajakumar, P.; Selvam, S. *Tetrahedron* **2007**, *63*, 8891-8901; (b) Rajakumar, P.; Padmanabhan, R. *Tetrahedron Lett.* **2010**, *51*, 1059-1063.
- 24. Hart, H.; Rajakumar, P. Tetrahedron 1995, 51, 1313-1336.
- 25. (a) Blazevic, N.; Kolbah, D.; Belin, B.; Sunjic, V.; Kajfez, F. Synthesis 1979, 161-165. (b) Bottini, A. T.; Vasu Dev.; Klinck, J. Organic Syntheses 1963, 43, 6-9. (c) Meyers, A. I.; Warmus, J. S.; Dilley, G. J. Organic Syntheses 1996, 73, 246-249. (d) Rajakumar, P.; Padmanabhan, R. Tetrahedron 2011, 67, 9669-9679.
- 26. Benesi, H. A.; Hildebrand, J. H. J. Am. Chem. Soc. 1949, 71, 2703-2707.