

Supplementary Material

A comparative study of solution phase as well as solvent free microwave assisted syntheses of 3-benzothiazole/isoxazole substituted 2-styryl-4(3H)-quinazolinones

**Chandresh L. Jagani,^a Satish F. Vanparia,^a Tarosh S. Patel,^a
Ritu B. Dixit,^b Bharat C. Dixit^{*a}**

^a V. P. & R. P. T. P. Science College, Affiliated to Sardar Patel University,
Vallabh Vidyanagar – 388120, Gujarat, India

^b Ashok & Rita Patel Institute of Integrated Studies and Research in Biotechnology and Allied
Sciences, New Vallabh Vidyanagar – 388121, Gujarat, India
E-mail: dixits20002003@yahoo.co.in




Figure 1. FT-IR spectrum of **2a**.




Figure 2. ^1H -NMR spectrum of 1a.




Figure 3. ^{13}C -NMR spectrum of **1a**.




Figure 4. Mass spectrum of 1a.




Figure 5. FT-IR spectrum of **1a**.




Figure 6. ¹H-NMR spectrum of **1o**.




Figure 7. ^{13}C -NMR (APT) spectrum of **1o**.




Figure 8. Mass spectrum of **1o**.




Figure 9. FT-IR spectrum of **1o**.