

Supplementary Material
Asymmetric Michael Addition of Acetone to β -Nitrostyrenes
Catalyzed by Novel Organocatalysts Derived from
D-Isomannide or L-Isoidide

Ling-Yan Chen,^{a,b} Stéphane Guillarme,^a Andrew Whiting^c and Christine Saluzzo^{a,*}

^a*IMMM, UMR CNRS 6283, Faculté des Sciences,
Université du Maine, Av. O. Messiaen, 72085 Le Mans Cedex 09, France*

^b*College of Chemistry and Chemical Engineering,
Shanghai University of Engineering Science, 201620, China*



^c*Department of Chemistry,
Durham University, Science Laboratories, South Road, Durham, DH1 3LE, UK*



E-mail: Christine.saluzzo@univ-lemans.fr



Table of contents



NMR data



¹ H NMR and ¹³ C NMR spectra of monothiourea 2	S2
¹ H NMR and ¹³ C NMR spectra of monothiourea 3	S3
¹ H NMR and ¹³ C NMR spectra of monothiourea 6	S4
¹ H NMR and ¹³ C NMR spectra of diimine 7	S5
¹ H NMR and ¹³ C NMR spectra of diimine 8	S6
¹ H NMR and ¹³ C NMR spectra of compound 10a	S7
¹ H NMR and ¹³ C NMR spectra of compound 10b	S8
¹ H NMR and ¹³ C NMR spectra of compound 10c	S9
¹ H NMR and ¹³ C NMR spectra of compound 10d	S10
¹ H NMR and ¹³ C NMR spectra of compound 10e	S11







¹H NMR spectrum of monothiurea 3**¹³C NMR spectrum of monothiurea 3**







¹H NMR spectrum of diimine 7**¹³C NMR spectrum of diimine 7**


¹H NMR spectrum of diimine 8**¹³C NMR spectrum of diimine 8**

¹H NMR spectrum of compound 10a**¹³C NMR spectrum of compound 10a**

¹H NMR spectrum of compound 10b**¹³C NMR spectrum of compound 10b**

¹H NMR spectrum of compound 10c**¹³C NMR spectrum of compound 10c**

¹H NMR spectrum of compound 10d**¹³C NMR spectrum of compound 10d**

¹H NMR spectrum of compound 10e**¹³C NMR spectrum of compound 10e**