

Supplementary Material

Protecting-group-free synthesis of GF109203X

Yong-Chen Gao,^b Yu-Hui Jia,^c Ting-Ting Li,^a Ze-Huan Hei,^a Feng-Ling Yang^b, Mu-Hua Huang,^{*a} and Yun-Jun Luo^{*a}

^a School of Materials Science and Engineering, Beijing Institute of Technology, Beijing, 100081, China

^b College of Chemistry and Molecular Engineering, Zhengzhou University, Zhengzhou, 450052 China

^c School of Chemistry, Beijing Institute of Technology, Beijing, 100081, China

E-mails: mhhuang@bit.edu.cn, yiluo@bit.edu.cn


Table of Contents

1. General Information	S2
2. ¹ H, ¹³ C-NMR, IR and HR-MS Spectrum	S3
3. Comparison of ¹ H-NMR Spectra of GF109203X	S22

1. General Information

All reagents were used as received from commercial sources without further purification or prepared as described in the literature. Tetrahydrofuran was distilled from sodium and benzophenone immediately before use. Reactions were stirred using Teflon-coated magnetic stir bars. Analytical TLC was performed with 0.20 mm silica gel 60F plates. Chromatographic purification of products was carried out by flash chromatography on silica gel (230-400 mesh). Melting points were determined using a electrothermal melting point apparatus or DSC. Infrared spectra were recorded on a Nicolet 8700 Fourier transform spectrometer. NMR spectra were measured in CDCl₃ (with TMS as internal standard) or DMSO-d₆ on a Bruker AV400 or Varian INOVA-400M (¹H at 400 MHz, ¹³C at 100 MHz) magnetic resonance spectrometer. Chemical shifts (δ) are reported in ppm, and coupling constants (*J*) are in Hz. The following abbreviations were used to explain the multiplicities: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet. High-resolution EI mass spectra (HR-EI-MS) were recorded on an GCT CA127 Micronass UK mass spectrometer.

2. ^1H , ^{13}C -NMR, IR and HR-MS Spectrum





—172.504


136,294
135,909
132,198
128,915
128,390
127,337
126,366
125,537
122,700
122,352
122,241
122,048
120,398
120,248
111,347
109,784
107,220
105,640



77,494
77,177
76,859


-46.963

32,054
30,127
29,757
29,633
29,490
29,363
27,010
22,836













C-D-0419-YCG-1-103-2
C-D-0419-YCG-1-103-2








Elemental Composition Report


Tolerance = 10.0 PPM / DBE: min = -1.5, max = 50.0
Isotope cluster parameters: Separation = 1.0 Abundance = 1.0%


Monoisotopic Mass, Odd and Even Electron Ions
180 formula(e) evaluated with 1 results within limits (up to 50 closest results for each mass)


Minimum:	80.00				-1.5				
Maximum:	100.00			200.0	10.0	50.0			
Mass	RA	Calc. Mass	mDa	PPM	DBE	Score	Formula		
495.2903	100.00	495.2886	1.7	3.5	16.0	1	C32 H37 N3 O2		



C-D-0418-YCG-1-103-1
C-D-0418-YCG-1-103-1








Elemental Composition Report


Tolerance = 10.0 PPM / DBE: min = -1.5, max = 50.0
Isotope cluster parameters: Separation = 1.0 Abundance = 1.0%


Monoisotopic Mass, Odd and Even Electron Ions



7 formula(e) evaluated with 1 results within limits (up to 50 closest results for each mass)

Minimum:	80.00								
Maximum:	100.00								
Mass	RA	Calc. Mass	mDa	PPM	DBE	Score	Formula		
663.4772	100.00	663.4764	0.8	1.2	16.0	1	C44 H61 N3 O2		











Elemental Composition Report


Tolerance = 10.0 PPM / DBE: min = -1.5, max = 50.0
Isotope cluster parameters: Separation = 1.0 Abundance = 1.0%

Monoisotopic Mass, Odd and Even Electron Ions
7 formula(e) evaluated with 1 results within limits (up to 50 closest results for each mass)

Minimum:	80.00					-1.5		
Maximum:	100.00		200.0	10.0	50.0			
Mass	RA	Calc. Mass	mDa	PPM	DBE	Score	Formula	
663.4776	100.00	663.4764	1.2	1.8	16.0	1	C44 H61 N3 O2	







3. Comparison of ^1H -NMR Spectra of GF109203X

Comparison of ^1H -NMR of our synthetic sample with the reported data of GF109203X (7)

Protons in 6	Our Synthetic Sample	GF109203X by Faul ^a	GF109203X by Toullec ^b
H _N ''	11.68 (s, 1H)	11.65 (bs, 1H)	11.66(s)
H _N	10.90 (s, 1H)	10.88 (s, 1H)	10.0-11.29 (bs)
H ₂ '	7.78 (d, $J = 2.4$ Hz, 1H)	7.75 (d, 1H, $J = 2.58$ Hz)	7.77 (bs)
H ₂ ''	7.70 (s, 1H)	7.66 (s, 1H)	7.68(s)
H ₄ ''	7.46 (d, $J = 8.0$ Hz, 1H)	7.37 (d, 1H, $J = 8.22$ Hz)	7.45 (d, $J = 8.2$ Hz)
H ₄ '	7.36 (d, $J = 8.8$ Hz, 1H)	7.16 (d, 1H, $J = 8.12$ Hz),	7.35 (d, $J = 8.1$ Hz)
H _{Ar} (3H)	7.03 (t, $J = 7.6$ Hz, 1H), 6.96 (t, $J = 7.6$ Hz, 2H), 6.86 (d, $J = 8.0$ Hz, 1H)	7.01-6.90 (m, 2H), 6.83 (d, 1H, $J = 7.96$ Hz)	7.01-6.83 (m)
H _{Ar} ' (3H)	6.70 (d, $J = 8.0$ Hz, 1H), 6.67 (t, $J = 7.6$ Hz, 1H), 6.57 (t, $J = 7.6$ Hz, 1H)	6.69-6.61 (m, 2H), 6.54 (t, 1H, $J = 7.54$ Hz)	6.72-6.56 (m)
CH ₂ (CH ₂) ₂ N(CH ₃) ₂	4.24 (t, $J = 6.4$ Hz, 2H)	4.21 (t, 2H, $J = 6.48$ Hz)	4.22 (t, $J = 6.5$ Hz),
CH ₂ CH ₂ N(CH ₃) ₂	2.09 (bs, 8H)	2.06-2.03 (bs, 8H),	2.06-2.03 (bs)
CH ₂ CH ₂ CH ₂	1.82 (t, $J = 6.8$ Hz, 2H)	1.81-1.77 (m, 2H);	1.83 (t, $J = 7.0$ Hz)

^a Faul, M. M.; Winneroski, L. L.; Krumrich, C. A. *J. Org. Chem.* **1998**, 63, 6053.

^b Toullec, D.; Pianetti, P.; Coste, H.; Bellevergue, P.; Grand-Perret, T.; Ajakane, M.; Baudet, V.; Boissin, P.; Boursier, E.; Loriolle, F. *J. Biol. Chem.* **1991**, 266, 15771.