

Supplementary Material

Regio - and stereoselective synthesis of the iminosugars – 4-substituted 1-benzylpiperidine-3,5-diols



Ivan S. Veselov*, Alexandr M. Petrenko, Dmitriy M. Mazur, and Galina V. Grishina*

Department of Chemistry, Moscow State University, 119991, Moscow, Russia


Email: ivan@aroma-os.ru, grishina@org.chem.msu.ru

Table of Contents



¹ H NMR spectrum (400 MHz, CD ₃ OD, DMSO d ₆) of 1-benzyl-4-benzylaminopiperidine-3,5-diol (10a)	S2
¹³ C NMR spectrum (100 MHz, CD ₃ OD, DMSO d ₆) of 1-benzyl-4-benzylaminopiperidine-3,5-diol (10a)	S3
¹ H NMR spectrum (400 MHz, CD ₃ OD) of 4-aminopiperidine-3,5-diol (11)	S4
¹³ C NMR spectrum (100 MHz, CD ₃ OD) of 4-aminopiperidine-3,5-diol (11)	S5




^1H NMR spectrum (400 MHz, CD_3OD , $\text{DMSO}-d_6$) of 1-benzyl-4-benzylaminopiperidine-3,5-diol (**10a**).



^{13}C NMR spectrum (100 MHz, CD_3OD , $\text{DMSO-}d_6$) of 1-benzyl-4-benzylaminopiperidine-3,5-diol (**10a**).



^1H NMR spectrum (400 MHz, CD_3OD) of 4-aminopiperidine-3,5-diol (**11**).



^{13}C NMR spectrum (100 MHz, CD_3OD) of 4-aminopiperidine-3,5-diol (**11**).