

Supplementary Material

An efficient synthesis of Nepetoidin B

Ming Yao ^{a*}, Jingjing Zhang ^{a, b}, Sen Yang ^a, Hangxing Xiong ^{a*}



^a *Jingchu University of Technology, Jingmen and 448000, P. R. China.*


^b *Wuhan Institute of Technology, Wuhan and 430205, P. R. China.*

Email: yaomingcep@jcut.edu.cn


Table of Contents

1. ¹ H NMR spectra of compounds 3	S2
2. ¹ H NMR spectra of compounds (E, E)-5	S2
3. ¹ H and ¹³ C NMR spectra of compounds (Z, E)-5	S3
4. ¹ H and ¹³ C NMR spectra of compounds (Z, E)-1/(E, E)-1	S4
5. ¹ H and ¹³ C NMR spectra of compounds (E, E)-1	S6


¹H NMR spectra of compounds **3**¹H NMR spectra of compounds **(E,E)-5**




¹H NMR spectra of compounds (Z, E)-5 (The EtOAc impurity in the spectra can be determined by ¹H-NMR spectroscopic analysis (signals at 4.12 (q, 3H), 2.05 (s, 2H) and 1.26 (t, 3H))




¹³C NMR spectra of compounds (Z, E)-5.




¹H NMR spectra of compounds (Z, E)-1/(E, E)-1 (acetone d₆, 400Hz)




¹³C NMR spectra of compounds (Z, E)-1/(E, E)-1 (acetone d₆, 100Hz)




¹H NMR spectra of compounds (Z, E)-1/(E, E)-1 (CD₃OD, 400Hz)




¹³C NMR spectra of compounds (Z, E)-1/(E, E)-1 (CD₃OD, 100Hz)




¹H NMR spectra of compounds (*E, E*)-1 (acetone d₆, 400 MHz)



¹³C NMR spectra of compounds (*E, E*)-1 (acetone d₆, 100 MHz)



¹H and ¹³C NMR spectra of compounds (*E, E*)-1 (DMSO *d*₆, 400Hz)



¹³C NMR spectra of compounds (*E, E*)-1 (DMSO *d*₆, 100Hz)