

Supplementary Material

The synthesis of 7,9-dimethoxy-3-propyl-3,4-dihydro-1*H*-benzo[*g*]isochromene-1,5,10-trione: A potential monomer for the synthesis of the natural product xylindein

Md. Firoj Hossain, Andreas Lemmerer,^{II} and Charles B. de Koning*

Molecular Sciences Institute, School of Chemistry, University of the Witwatersrand, PO Wits 2050, Johannesburg, South Africa

Email: Charles.deKoning@wits.ac.za

Table of Contents

¹ H and ¹³ C NMR spectra of ethyl-3-allyl-4-hydroxy-1,6,8-trimethoxynaphthalene-2-naphthoate (12)	S2
¹ H and ¹³ C NMR spectra of ethyl-3-allyl-4-(benzyloxy)-1,6,8-trimethoxy-2-naphthoate (13)	S3
¹ H and ¹³ C NMR spectra of ethyl-4-(benzyloxy)-1,6,8-trimethoxy-3-(4-oxopent-2-enyl)-2-naphthoate (14)	S4
¹ H and ¹³ C NMR spectra of 5-(benzyloxy)-7,9,10-trimethoxy-3-(2-oxopropyl)-3,4-dihydro-1 <i>H</i> -benzo[<i>g</i>]isochromen-1-one (15)	S5
¹ H and ¹³ C NMR spectra of 1-[10-(benzyloxy)-1-hydroxy-6,8,9-trimethoxyanthracen-2-yl]ethanone (16)	S6
¹ H and ¹³ C NMR spectra of 5-(benzyloxy)-10-hydroxy-7,9-dimethoxy-3-(2-methyl-1,3-dithiolan-2-yl)methyl-3,4-dihydro-1 <i>H</i> -benzo[<i>g</i>]isochromen-1-one (17a)	S7
¹ H and ¹³ C NMR spectra of 5-(benzyloxy)-7,9,10-trimethoxy-3-(2-methyl-1,3-dithiolan-2-yl)methyl-3,4-dihydro-1 <i>H</i> -benzo[<i>g</i>]isochromen-1-one (17)	S8
¹ H and ¹³ C NMR spectra of 5-hydroxy-7,9-dimethoxy-3-propyl-3,4-dihydro-1 <i>H</i> -benzo[<i>g</i>]isochromen-1-one (18)	S9
¹ H and ¹³ C NMR spectra of 7,9-dimethoxy-3-propyl-3,4-dihydro-1 <i>H</i> -benzo[<i>g</i>]isochromen-1,5,10-trione (10)	S10





Figure S1. ¹H and ¹³C NMR spectra (CDCl₃) of ethyl-3-allyl-4-hydroxy-1,6,8-trimethoxynaphthalene-2-naphthoate (**12**).






Figure S3. ¹H and ¹³C NMR spectra (CDCl₃) of ethyl-4-(benzyloxy)-1,6,8-trimethoxy-3-(4-oxopent-2-enyl)-2-naphthoate (**14**).




Figure S4. ¹H and ¹³C NMR spectra (CDCl₃) of 5-(benzyloxy)-7,9,10-trimethoxy-3-(2-oxopropyl)-3,4-dihydro-1H-benzo[g]isochromen-1-one (**15**).




Figure S5. ¹H and ¹³C NMR spectra (CDCl₃) of 1-[10-(benzyloxy)-1-hydroxy-6,8,9-trimethoxyanthracen-2-yl]ethanone (**16**).




Figure S6. ¹H and ¹³C NMR spectra (CDCl₃) of 5-(benzyloxy)-10-hydroxy-7,9-dimethoxy-3-(2-methyl-1,3-dithiolan-2-yl)methyl-3,4-dihydro-1H-benzo[g]isochromen-1-one (17a).





Figure S7. ¹H and ¹³C NMR spectra (CDCl₃) of 5-(benzyloxy)-7,9,10-trimethoxy-3-(2-methyl-1,3-dithiolan-2-yl)methyl-3,4-dihydro-1H-benzo[g]-isochromen-1-one (**17**).






Figure S9. ¹H and ¹³C NMR spectra (CDCl₃) of 7,9-dimethoxy-3-propyl-3,4-dihydro-1H-benzo[g]isochromen-1,5,10-trione (10).