

Supplementary Material

Synthesis and application of a fluorescent “turn-off” triazolyl-coumarin-based fluorescent chemosensor for the sensing of Fe³⁺ ions in aqueous solutions

Neliswa Mama* and Aidan Battison

Department of Chemistry, Nelson Mandela University, P.O Box 77000, Port Elizabeth, 6031, South Africa

Email: neliswa.mama@mandela.ac.za

Table of Contents

S1. FT-IR spectrum of “click” product S1	S2
S2. FT-IR spectrum of “click” product S2	S2
S3. ¹ H NMR spectrum of sensor S1 in DMSO-d ₆	S3
S4. ¹ H NMR spectrum of sensor S2 in DMSO-d ₆	S3
S5. ¹ H NMR spectrum of 1-(prop-2-ynyloxy)benzene derivative 2 in CDCl ₃	S4
S6. ¹³ C NMR spectrum of sensor S1 in DMSO-d ₆	S4
S7. ¹³ C NMR spectrum of sensor S2 in DMSO-d ₆	S5
S8. Time-delay studies of S1 with a single aliquot of Fe ³⁺	S5
S9. Benesi-Hildebrand plot of S1 in acetonitrile.....	S6




Figure S1: FT-IR spectrum of “click” product **S1**.




Figure S2: FT-IR spectrum of “click” product **S2**.




Figure S3: ¹H NMR spectrum of sensor **S1** in DMSO-d₆




Figure S4: ¹H NMR spectrum of sensor **S2** in DMSO-d₆




Figure S5: ¹H NMR spectrum of 1-(prop-2-ynyloxy)benzene derivative **2** in CDCl₃.




Figure S6: ¹³C NMR spectrum of sensor **S1** in DMSO-d₆.




Figure S7: ¹³C NMR spectrum of sensor **S2** in DMSO-d₆.




Figure S8: Time-delay studies of **S1** with a single aliquot of Fe³⁺.




Figure S9: Benesi-Hildebrand plot of **S1** in acetonitrile.