

Synthesis and structural elucidation of 2,3-dimethylnaphthazarin ester derivatives

Mehmet F. Saglam*

Department of Chemistry, Faculty of Science, Gebze Technical University, 41400, Kocaeli, Turkey Email: <u>mfsaglam@gtu.edu.tr</u>

Received 12-30-2022

Accepted Manuscript 02-14-2023

Published on line 02-26-2023

Abstract

A simple and efficient synthesis of bis-ester substituted 2,3-dimethylnaphthazarins is described. The preparation of 5-*O*,8-*O*-bis-ester derivatives of 2,3-dimethylnaphthazarin was achieved via double esterification reaction of 5,8-dihydroxy-2,3-dimethylnaphthazarin with a range of acyl chlorides under mild reaction conditions. Additionally, 5-*O*- mono ester and unsymmetrical 5-*O*,8-*O*-bis-ester derivatives of 2,3-dimethylnaphthazarins were successfully synthesized. The identities of new compounds were confirmed by ¹H NMR, ¹³C NMR, FTIR, HRMS and X-ray single-crystal diffraction techniques.



Keywords: Naphthazarins, synthesis, esterification, structure elucidation

Introduction

Naphthazarin (5,8-dihydroxy-1,4-naphthoquinone) and congeners constitute an important class of compounds which have been isolated from natural sources and display a wide range of pharmacological activities.¹⁻³ Synthetic analogues of the naphthazarins also possess potent and often valuable biological activities such as antimicrobial, antibacterial, antiviral, anticancer, anti-inflammatory, analgesic and antithrombotic effects.⁴⁻⁶ Well known naphthazarin-based compounds, such as Shikonin and its enantiomer Alkannin (Figure 1) have been isolated from in the roots of *Lithospermum erythrorhizon* and *Arnebia euchroma*.^{5,7} Among the naphthazarin derivatives, 2,3-dimethylnaphthazarin (DMN) is of significant synthetic interest due to its biological activity.⁸ In particular, bis-benzoyl ester dimethylnapthazarin derivatives have been reported to possess borderline inhibitory activity against leukemia P388 cells.⁸ Apart from the pharmaceutical importance, DMN-based compounds have been recognized as being of value and their synthesis has thus been targeted by organic chemists.



Figure 1. Naphthazarin and its biologically important derivatives.

Limited synthetic approaches have been reported for the synthesis of DMN and its derivatives. One of the oldest methods was reported by Rodriguez in 1984 which involved a multi-step procedure including an initial cycloaddition between 2,3-dimethyl-1,3-butadiene and *p*-benzoquinone. This step was followed by acetylation and chromium trioxide oxidation of the resulting adducts and subsequent hydrolysis of the resultant naphthazarin diacetate.⁹ In a similar manner, DMN has also been prepared from 1,3-butadiene and 2,3-dimethylquinone.⁹ The most efficient, and more recent synthesis of DMN has been achieved in a one-pot process by the direct reaction of 2,3-dimethylmaleic anhydride and 1,4-dimethoxybenzene, in the presence of aluminum chloride and sodium chloride.¹⁰ The main advantage of this latter method is that the DMN could be synthesized in relatively high yield from inexpensive commercial chemicals.

In conjunction with the synthetic efforts focused on naphthazarin, the presence of the hydroxyl groups, located at the C5 and C8 positions on the naphthazarin aromatic ring, should allow the synthesis of a range of targeted mono- and bis- ester compounds. It is known that a range of ester derivatives have been prepared from Shikonin and Alkannin, wherein the secondary alcohol functionality located on the side chain were converted to the corresponding *O*-ester.¹¹⁻¹² However, (selective) derivatization of the parent DMN has rarely been studied in the literature,¹⁰ presumably due to the harsh reaction conditions and multi-step processes involved in the previously reported synthetic routes. The current study presents the development of a general method for the synthesis of bis-ester 2,3-dimethylnaphthazarin derivatives from reaction of DMN with acyl chlorides under mild reaction conditions. The reaction is facilitated by the phenoxy groups located at the C5 and C8 positions and yields both mono-ester and unsymmetrical bis-ester derivatives of the parent 2,3-dimethylnaphthazarin.

Results and Discussion

In earlier studies, the synthesis of bis-acetyl and bis-benzoyl ester derivatives of 2,3-dimethylnapthazarin 5 and 12 were prepared via multi-step reactions, requiring harsh reaction conditions and long reaction times. For example, Cheng and coworkers reported the synthesis of bis-acetyl-2,3-dimethylnaphthazarin 5 in three steps starting from building the naphthoquinone platform via a Diels-Alder reaction of 2,3-dimethyl-1,3-butadiene and p-benzoguinone, followed by acetylation and oxidation with chromium trioxide.⁸ The yield of the reaction was mentioned as extremely low and no characterization data was reported for the bis-acetate product 5. Using the same synthetic route, the group also reported the synthesis of bis-benzoyl derivative 12 in three steps using benzoyl chloride as an acylation reagent. The overall yield for this reaction sequence was likewise very low, as the yield from the final oxidation was only 14%. In addition, the total reaction time was reported to be 32 hours. The detailed characterization data for derivative **12** was also not reported. In 1986, Rodriguez and coworkers reported the synthesis of the bis-acetyl ester of 2,3-dimethylnaphthazarin 5 following the same reaction sequence.¹³ After building the naphthoquinone structure using a Diels-Alder reaction, treatment of the resulting adduct with acetic anhydride in acetic acid under reflux for 3 hours and subsequent oxidation with chromium trioxide generated product 5. The final oxidation step provided bis-acetyl-2,3-dimethylnaphthazarin 5 in 35% yield and ¹H NMR characterization data of product **5** was provided. To best of our knowledge, there is no other study in the literature for the synthesis of symmetrical bis-ester derivatives of the parent 2,3dimethylnapthazarin. In addition to this, the preparation of mono-ester or unsymmetrical bis-ester derivatives of DMN have not been reported thus far.

In order to bridge this gap, DMN **3** was synthesized by following a literature procedure¹⁰ involving a double Friedel-Craft acylation reaction between 1,4-dimethoxybenzene **1** and 2,3-dimethylmaleic anhydride **2** in the presence of molten form of aluminum chloride-sodium chloride, providing DMN **3** in an isolated yield of 37% (Scheme 1).



Scheme 1. Synthesis of 2,3-dimethylnaphthazarin.

With DMN **3** in hand, reaction conditions for its double esterification were investigated using acetyl chloride **4** as an acylation reagent. Initially, one equivalent of naphthazarin **3** was treated with three equivalents of base at room temperature, followed by four equivalents of acetyl chloride and the reaction progress was monitored by TLC analysis. When potassium carbonate or pyridine was used as the base, only a trace amount of the target product **5** was isolated. When the base was changed to triethylamine, the reaction gave the targeted bis-ester **5** in 28% yield along with its mono-ester, 5-*O*-acetyl-2,3-dimethylnaphthazarin **6**, in 4% yield. Following optimization, it was found that the treatment of naphthazarin **3** with three equivalents of triethylamine at room temperature for 30 minutes, followed by the addition of acetyl chloride **4** at 0 °C and subsequent warming to room temperature provided the highest yield of the target product **5** in 45% yield, along

with the 5-O-acetyl-mono-ester **6** in 3% yield (Scheme 2). The structures of products **5** and **6** were elucidated by a detailed one- and two-dimensional NMR spectroscopy investigation, as well as being fully characterized using FTIR, HRMS and finally confirmed by single crystal X-ray analysis (Figure 2).





The ¹H NMR spectrum of the bis-ester **5** at room temperature in CDCl₃ is consistent with that reported in the literature.¹³ The most important evidence for successful double esterification of **3** is the disappearance of both of the phenolic O*H* signals which initially appear as a singlet at δ 12.63 ppm due to intramolecular hydrogen-bonding interaction between the carbonyl oxygen of quinonoid ring and the proton of phenolic hydroxyl group proton in the aromatic ring of **3**. The appearance of a new singlet signal at δ 2.44 ppm in the proton NMR spectrum of **5** was associated with the protons of the two acetyl ester C*H*₃ group. In the ¹³C NMR spectrum of the bis-product **5**, the appearance of new signals in the high field region was associated with acetyl methyl *C*H₃ carbons (δ 21.2 ppm), while additional signals in the low field region were associated with the acetyl ester carbonyl *C*=O carbons (δ 169.5 ppm), proving that the double esterification of **3** has taken place. Highresolution mass spectrometry indicated that the molecular ion of product **5** has a mass to charge ratio of 325.0676, consistent with the molecular formula C₁₆H₁₄O₆ [M+Na]⁺. The structure of bis-ester **5** was further confirmed by single crystal X-ray crystallography analysis (Figure 2).

The new minor mono-ester **6** was also fully characterized. HRMS showed that the molecular ion of product **6** has a mass to charge ratio of 260.0686, consistent with the molecular formula $C_{14}H_{12}O_5$ [M]⁺. Finally, an analytical sample of mono-ester **6** was successfully recrystallized and its structure was also confirmed by single crystal X-ray (Figure 2).



Figure 2. X-ray structures of 5-*O*,8-*O*-diacetyl-bis-ester **5** and 5-*O*-acetyl-mono-ester **6** derivatives of 2,3-dimethylnaphthazarin.

The structure of mono-ester 6 was elucidated with further NMR experiments, namely APT NMR, 2dimensional ¹H -¹H (COSY), and ¹H-¹³C (HMQC, HMBC) correlation experiments in CDCl₃ at room temperature. In the ¹H NMR spectrum of **6**, only one phenolic O**H** signal was observed at δ 12.62 ppm as a singlet. This signal proved that the single esterification took place. As mono-ester **6** is not symmetrical, unlike the starting material DMN 3 and bis-ester 5, both of the methyl group CH_3 protons on the quinonoid ring give different proton resonances as singlets at δ 2.15 and 2.12 ppm. The ¹³C NMR spectrum of mono-ester **6** showed the methyl **C**H₃ carbons on quinonoid ring at different chemical shifts of δ 13.2 and 12.3 ppm. The signals appearing at δ 21.2 ppm and 169.8 ppm associated with the acetyl ester methyl carbon CH_3 and carbonyl carbon C=0, respectively. It is worth to note that while the carbonyl carbon signal (C1) on the quinonoid ring shifted to downfield region and appears at δ 189.9 ppm, the other carbonyl carbon signal (C4) resonates at δ 182.9 ppm, which is further upfield when compared to the starting material **3** (δ 186.5 ppm) (Table 1). It shows that the carbonyl oxygen C=O at C1 is intramolecularly hydrogen bonded with the phenolic OH proton of the aromatic ring, whereas the C4 carbonyl oxygen is not. Additionally, carbon atoms attached to the phenolic hydroxyl group showed significant chemical shift differences. For example, the phenoxy group attached to carbon C8 of mono-ester 6 resonated at δ 159.9 ppm, similar to DMN **3** (Table 1). However, once the phenolic hydroxyl group was converted into the corresponding ester derivative 5, the carbon atom (C5) signal at the aromatic ring appeared in the far upfield region at δ 142.6 ppm, as compared to DMN **3** (δ 158.6) (Table 1). Another important chemical shift difference was observed at the fused carbon atoms such as C10, which is next to the acetyl substituted carbon (C5) on the aromatic ring, which resonated downfield at δ 122.1 ppm, whereas carbon C9, neighboring to phenolic hydroxyl substituted carbon (C8), resonated at δ 114.8 ppm. On the contrary, the fused carbon signals of bis-ester **5** appeared in the far downfield region at δ 124.7 ppm (Table 1).

In order to extend this study to generate a variety of new 2,3-dimethylnaphthazarin bis-ester derivatives, various commercially available aromatic acyl , such as benzoyl, *p*-methyl-, *p*-bromo-, *p*-methoxy-, and *p*-nitrobenzoyl chlorides (**7-11**), were also used (Table 2). The esterification reaction was performed using the optimum reaction conditions as described earlier. The reaction progress was monitored by TLC analysis and the reaction was worked up once no further change was observed on the TLC. It was found that the reaction with *p*-bromobenzoyl chloride **9** was completed in 1.5 hours, whereas the reaction with benzoyl chloride **7** took 22 hours; both of them gave bis-ester **14** and **12** in 30% and 23% yield, respectively, as sole products (Table 2). The reaction with *p*-methylbenzoyl chloride **8** resulted in bis-adduct **13** as major product in 30% yield, along with mono-ester **17** in 5% yield, while the reaction with *p*-nitrobenzoyl chloride **11** gave bis-ester **16** and mono-ester **19** products, 21% and 27%, respectively. In contrast, the reaction between 2,3-dimethylnapthazarin **3** and *p*-methoxybenzoyl chloride **10** resulted in mono-ester **18** in 73% yield as a major product along with bis-ester **15** in 13% yield as the minor product.

Table 1. ¹H and ¹³C NMR chemical shifts of DMN 3, 5-O-acetyl mono-ester 6, and 5-O,8-O-diacetyl bis-ester 5



Position	DMN 3		Mono-ester 6		Bis-ester 5	
	$\delta_{ extsf{H}}$ / ppm	$\delta_{ m C}$ / ppm	$\delta_{ extsf{H}}$ / ppm	$\delta_{ m C}$ / ppm	$\delta_{ extsf{H}}$ / ppm	$\delta_{ m C}$ / ppm
1	-	186.5	-	189.9	-	183.2
2	-	144.5	-	142.3	-	143.5
3	-	144.5	-	145.8	-	143.5
4	-	186.5	-	182.9	-	183.2
5	12.63 (OH)	158.6	-	142.6	-	147.3
6	7.20	129.4	7.24	132.6	7.32	130.3
7	7.20	129.4	7.23	125.5	7.32	130.3
8	12.63 (OH)	158.6	12.62 (OH)	159.9	-	147.3
9	-	111.7	-	114.8	-	124.7
10	-	111.7	-	122.1	-	124.7
11	2.20	12.6	2.15	12.3	2.08	13.0
12	2.20	12.6	2.12	13.2	2.08	13.0
13	NA	-	-	169.8	-	169.5
14	NA	-	2.42	21.2	2.44	21.2
13′	NA	-	NA	-	-	169.5
14′	NA	-	NA	-	2.44	21.2

¹H NMR (in CDCl₃, 500 MHz), ¹³C NMR (in CDCl₃, 125 MHz).

Table 2. Synthesis of symmetrical bis-ester derivatives of 2,3-dimethylnaphthazarin



Following the successful synthesis of symmetrical bis-ester derivatives of DMN (**5**,**12-16**), the synthesis of an *unsymmetrical* bis-ester of DMN was examined. By using similar reaction conditions, that is treating one equivalent of DMN **3** with one equivalent of acetyl chloride **4** in THF, 5-*O*-acetyl-mono-ester **6** was generated in 36% yield, bis-ester **5** in 29% yield, while 3% of unreacted DMN **3** was recovered (Scheme 3). One equivalent of mono-ester **6** was then treated with one and a half equivalents of benzoyl chloride **7** at room temperature, in the presence of one and a half equivalents of triethylamine and THF, to result in the *unsymmetrical* bis-ester **20** in 17% yield (Scheme 3). This product **20** was fully characterized using spectroscopic methods and its structure was also confirmed by single crystal x-ray crystallography.





Conclusions

In summary, a variety of new ester derivatives of 2,3-dimethylnaphthazarin was achieved under mild reaction conditions in one step. These new dimethylnaphthazarin structures have considerable potential for development in areas including medicinal chemistry. Additionally, mono-ester derivatives create an opportunity to generate a library of *unsymmetrical* bis-substituted naphthazarins, and this work is currently in progress.

Experimental Section

General Information. Reactions were performed under a positive pressure of nitrogen or argon in oven or flame dried glassware, unless otherwise specified. All reagents and solvents were obtained from commercial sources and appropriately purified, if necessary. Anhydrous THF was dried over sodium wire and distilled from sodium benzophenone ketyl. ¹H and ¹³C NMR spectra were recorded at 298 K on a Varian 500 MHz NMR spectrometer. The deuterated solvent (deutero chloroform (CDCl₃) for NMR spectroscopy was obtained from Merck. The central line of the residual chloroform (CHCl₃ in CDCl₃) (δ = 7.26 ppm) and triplet (δ = 77.10 ppm) was used for ¹H NMR and ¹³C NMR spectra as an internal reference, respectively. The following abbreviations and combinations thereof are used to describe ¹H NMR spectra multiplicities: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, br. = broad. Assignment of proton signals was assisted by COSY experiment where necessary. Assignment of carbon signals was assisted by APT, HMBC, and HSQC experiment where necessary. IR spectra were recorded on between 4000 and 650 cm⁻¹ using a Perkin Elmer Spectrum 100 FT-IR spectrometer with an attenuated total reflection (ATR) accessory. Mass spectra were recorded on a Thermo ORBITRAP Q-EXACTIVE mass spectrometer (Bremen, Germany) equipped with a Troyasil (Istanbul Turkiye) C18 column (150 x 3 mm i.e., 3 µm particle size) for measurements of HRMS. Melting points were measured using a Stuart SMP10

melting point apparatus. Analytical TLC was performed with Merck silica gel plates, precoated with silica gel 60 F254 (0.2 mm) on aluminium sheets and visualized using UV fluorescence (λ max = 254 nm). Flash column chromatography employed Merck Kieselgel 60 (230–400 mesh) silica gel.

Experimental procedures and characterization data

5,8-Dihydroxy-2,3-dimethylnaphthalene-1,4-dione (DMN) (3). This compound **3** was prepared following the literature procedure.¹⁰ A two necked 100 mL reaction flask was charged with dry AlCl₃ (6.0 g, 45 mmol) and NaCl (1.35 g, 23 mmol) under nitrogen gas atmosphere. The mixture stirred and heated to 180 °C to provide a melt. To this was carefully added 1,4-dimethoxybenzene **1** (0.69 g, 5.0 mmol) and 2,3-dimethylmaleic anhydride **2** (1.9 g, 15 mmol) portion-wise (4 portions) which resulted in a dark red mixture. The reaction mixture was heated to 200 °C and maintained at this temperature for 5 minutes. After gas evolution ceased the reaction mixture was cooled to 100 °C, after which a ice-cold solution of 10% HCl (60 mL) was carefully added portion-wise (10 mL/each portion) to afford a maroon-coloured suspension. The reaction mixture was then allowed to cool to room temperature and stirred for overnight. The resulting mixture was extracted with DCM (5 x 20 mL) and the organic phases were dried over MgSO₄, filtered and concentrated under reduced pressure to afford the crude product as a red solid (1.84 g). Purification by flash column chromatography (SiO₂, acetone:hexane (15:85)) gave the title compound **3** (0.41 g, 1.87 mmol, 37% yield) as a red shiny solid, as well as recovered 2,3-dimethylmaleic anhydride **2** (0.71 g). R_f 0.43 (acetone:hexane (10:90)); mp 166-168 °C.¹³ The ¹H NMR spectroscopy data for 2,3-dimethylnaphthazarin (DMN) **3** was in agreement with that reported in the literature.¹⁰ ¹H NMR (500 MHz, CDCl₃) δ 12.63 (s, 2H), 7.20 (s, 2H), 2.20 (s, 6H) ppm.

General Procedure-1. Double esterification reactions of 2,3-dimethylnaphthazarin (DMN). To a stirred solution of 2,3-dimethylnaphthazarin (DMN) **3** in dry THF under argon atmosphere at room temperature, dry three equivalents of Et_3N was added dropwise and the dark red colour reaction mixture was stirred for 30 minutes at 25 °C in the dark. The reaction mixture was then cooled to 0 °C and stirred for 10 minutes before four equivalents of an acyl chloride was added. The resulting reaction mixture was then warmed to room temperature and stirred for **1.5-22** hours. The reaction solvent was then removed by rotary evaporator and the resulted residue subjected to quick filtration using SiO₂. The solvent of the collected fractions were then concentrated and subjected to column chromatography to give the target product.

6,7-Dimethyl-5,8-dioxo-5,8-dihydronaphthalene-1,4-diyl diacetate (5). The bis-acetylester DMN **5** was synthesized following general procedure-1 using DMN **3** (50 mg, 0.23 mmol) and acetyl chloride **4** (65 μ L, 0.92 mmol) in dry THF (5.0 mL) and dry Et₃N (96 μ l, 0.69 mmol). The reaction mixture was stirred at 25 °C for 3 hours. The resulting reaction solvent was then removed by rotary evaporator and the crude material was subjected to column chromatography (EtOAc:Hexane (20:80)) to give the target bis-ester compound **5** (31 mg, 45%) as orange colour solid, along with the mono-ester by-product **6** (2 mg, 3%) as yellow coloured solid. The ¹H NMR data for bis-acetyl-2,3-dimethylnaphthazarin **5** was in agreement with that reported in the literature.¹³ An analytical sample of **5** was obtained by recrystallization from dichloromethane/hexane to give yellow color needles. R_f 0.24 (in ethyl acetate:hexane (20:80)); mp 185-187 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.32 (s, 2H), 2.44 (s, 6H), 2.08 (s, 6H) ppm; ¹³C NMR (125 MHz, CDCl₃) 183.2, 169.5, 147.3, 143.5, 130.3, 124.7, 21.2, 13.0 ppm; FT-IR: \bar{v}_{max} 3081, 3007, 1753, 1647, 1589, 1466, 1369, 1263, 1182, 937, 764 cm⁻¹; HRMS: calc for C₁₆H₁₄O₆ [M+Na]⁺: 325.0688; found 325.0676.

4-Hydroxy-6,7-dimethyl-5,8-dioxo-5,8-dihydronaphthalen-1-yl acetate (6). R_f 0.37 (in ethyl acetate:hexane (20:80)); mp 143-145 °C; ¹H NMR (500 MHz, CDCl₃) δ 12.62 (s, 1H), 7.24 (s, 1H), 7.23 (s, 1H), 2.42 (s, 3H), 2.15 (s, 3H), 2.12 (s, 3H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ 189.9, 182.9, 169.8, 159.9, 145.8, 142.6, 142.3, 132.6, 125.5,

122.1, 114.8, 21.2, 13.2, 12.3 ppm; FT-IR: $\bar{\nu}_{max}$ 2920, 2850, 1757, 1652, 1634, 1618, 1461, 1435, 1359, 1280, 1186, 1011, 786 cm⁻¹; HRMS: calc for C₁₄H₁₂O₅ [M]⁺: 260.0685; found 260.0686.

6,7-Dimethyl-5,8-dioxo-5,8-dihydronaphthalene-1,4-diyl dibenzoate (12). The bis-benzoylester DMN **12** was synthesized following general procedure-1 using DMN **3** (50 mg, 0.23 mmol) and benzoyl chloride **7** (0.27 mL, 2.3 mmol) in dry THF (5.0 mL) and dry Et₃N (96 µl, 0.69 mmol). The reaction mixture was stirred at 25 °C for 22 hours. The resulting reaction solvent was then removed by rotary evaporator and the crude material was subjected to column chromatography (EtOAc:Hexane (20:80)) to give the compound **12** (22 mg, 23%) as an orange coloured solid. An analytical sample of **12** was obtained by recrystallization from dichloromethane/hexane to give orange color needles. R_f 0.67 (in dichloromethane); mp 274-276 °C;^{8 1}H NMR (500 MHz, CDCl₃) δ 8.26 (dd, *J* 11.3, 10.0 Hz, 4H), 7.71 – 7.65 (m, 2H), 7.60 – 7.53 (m, 4H), 7.51 (s, 2H), 2.04 (s, 6H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ 183.2, 165.2, 147.7, 143.5, 133.8, 130.6, 130.5, 129.4, 128.8, 125.1, 13.0 ppm; FT-IR: $\bar{\nu}_{max}$ 2919, 1733, 1650, 1636, 1584, 1491, 1466, 1450, 1418, 1262, 1219, 1064, 702 cm⁻¹; HRMS: calc for C₂₆H₁₈O₆Na [M+Na]⁺: 449.1001; found 449.0988.

6,7-Dimethyl-5,8-dioxo-5,8-dihydronaphthalene-1,4-diyl bis(4-methylbenzoate) (13). The bis-*p*-methylbenzoylester DMN **13** was synthesized following general procedure-1 using DMN **3** (50 mg, 0.23 mmol) and *p*-methylbenzoyl chloride **8** (0.12 mL, 0.92 mmol) in dry THF (2.0 mL) and dry Et₃N (96 µl, 0.69 mmol). The reaction mixture was stirred at 25 °C for 1.5 hours. The resulting reaction solvent was then removed by rotary evaporator and the crude material was subjected to column chromatography (EtOAc:Hexane (20:80)) to give the compound **13** (31 mg, 30%) as yellow coloured solid along with by product mono-ester **17** (4 mg, 5%) as an orange-coloured solid. R_f 0.64 ((in dichloromethane:ethanol (99:1)); mp 265-267 °C; ¹H NMR (500 MHz, CDCl₃) δ ppm; NMR (500 MHz,) δ 8.15 (d, *J* 8.0 Hz, 4H), 7.49 (s, 2H), 7.36 (d, *J* 8.0 Hz, 4H), 2.48 (s, 6H), 2.03 (s, 6H); ¹³C NMR (125 MHz, CDCl₃) δ ppm; ¹³C NMR (126 MHz, CDCl₃) δ 183.15, 165.21, 147.67, 144.68, 143.48, 130.61, 130.49, 129.48, 126.69, 125.10, 21.91, 12.99; FT-IR: \bar{v}_{max} 2916, 1752, 1743, 1658, 1578, 1271, 1223, 1178, 1077, 734 cm⁻¹; HRMS: calc for C₂₈H₂₂O₄Na [M+Na]⁺: 477.1314; found 477.1303.

4-Hydroxy-6,7-dimethyl-5,8-dioxo-5,8-dihydronaphthalen-1-yl 4-methylbenzoate (17). R_f 0.37 (in ethylacetate:hexane (20:80)); mp 249-251 °C; ¹H NMR (500 MHz, CDCl₃) δ 12.68 (s, 1H), 8.13 (d, *J* 8.0 Hz, 2H), 7.37 – 7.31 (m, 3H), 7.31 – 7.27 (m, 1H), 2.47 (s, 3H), 2.14 (s, 3H), 2.07 (s, 3H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ 190.0, 182.7, 165.4, 159.9, 146.0, 144.6, 142.8, 142.2, 132.9, 130.5, 129.5, 126.8, 125.4, 122.5, 114.9, 21.9, 13.3, 12.3 ppm; FT-IR: \bar{v}_{max} 2919, 2850, 1727, 1687, 1614, 1469, 1225, 1086, 1059, 737 cm⁻¹; HRMS: calc for C₂₀H₁₆O₅Na [M+Na]⁺: 359.0895; found 359.0884.

6,7-Dimethyl-5,8-dioxo-5,8-dihydronaphthalene-1,4-diyl bis(4-bromobenzoate) (14). The bis-*p*-bromobenzoylester DMN **14** was synthesized following general procedure-1 using DMN **3** (50 mg, 0.23 mmol) and *p*-bromobenzoyl chloride **9** (0.20 g, 0.92 mmol) in dry THF (2.0 mL) and dry Et₃N (96 µl, 0.69 mmol). The reaction mixture was stirred at 25 °C for 1.5 hours. The resulting reaction mixture subjected a quick filtration with silica gel and combined fraction solvent was then removed by rotary evaporator. The crude material was subjected to column chromatography eluting with dichloromethane to give the compound **14** (40 mg, 30%) as orange-coloured solid. R_f 0.59 (in dichloromethane); mp 293-295 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.12 (d, *J* 8.3 Hz, 4H), 7.71 (d, *J* 8.4 Hz, 4H), 7.50 (s, 2H), 2.03 (s, 6H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ 183.1, 164.5, 147.5, 143.6, 132.2, 132.1, 130.4, 129.2, 128.3, 125.0, 13.0 ppm; FT-IR: $\bar{\nu}_{max}$ 1743, 1657, 1584, 1465, 1377, 1222, 1078, 1010, 843, 741 cm⁻¹; HRMS: calc for C₂₆H₁₆⁸¹Br₂O₆ [M]⁺: 608.9191; found 608.9166, calc for C₂₆H₁₆⁸¹Br⁷⁹BrO₆ [M]⁺: 604.9211; found 604.9207.

6,7-Dimethyl-5,8-dioxo-5,8-dihydronaphthalene-1,4-diyl bis(4-methoxybenzoate) (15). The bis-p-methoxybenzoylester DMN 15 was synthesized following general procedure-1 using DMN 3 (50 mg, 0.23 mmol) and p-methoxybenzoyl chloride 10 (0.15 g, 0.92 mmol) in dry THF (2.0 mL) and dry Et₃N (96 μ l, 0.69 mmol). The

reaction mixture was stirred at 25 °C for 2.5 hours. The resulting reaction mixture subjected a quick filtration with silica gel and combined fraction solvent was then removed by rotary evaporator. The crude material was subjected to column chromatography (dichloromethane:hexane (90:10)) to give the target bis-product **15** (15 mg, 13%) as yellow-coloured solid, along with mono-product **18** (59 mg, 73%) as by-product as an orange-coloured solid. R_f 0.30 (in dichloromethane:hexane (90:10)); mp 240-242 °C. ¹H NMR (500 MHz, CDCl₃) δ 8.22 (d, *J* 8.7 Hz, 4H), 7.48 (s, 2H), 7.03 (d, *J* 8.7 Hz, 4H), 3.91 (s, 6H), 2.03 (s, 6H) ppm; ¹³C NMR (126 MHz, CDCl₃) δ 183.2, 164.8, 164.1, 147.7, 143.5, 132.7, 130.5, 125.1, 121.8, 114.1, 55.6, 13.0 ppm; FT-IR: \bar{v}_{max} 1752, 1660, 1608, 1578, 1511, 1224, 1167, 1080, 840, 682 cm⁻¹; HRMS: calc for C₂₈H₂₂O₈ [M+Na]⁺: 509.1212; found 509.1203. **4-Hydroxy-6,7-dimethyl-5,8-dioxo-5,8-dihydronaphthalen-1-yl 4-methoxybenzoate** (**18**): R_f 0.49 (in dichloromethane:hexane (90:10)); mp 167-169 °C; ¹H NMR (500 MHz, CDCl₃) δ 12.67 (s, 1H), 8.19 (d, *J* 8.7 Hz, 2H), 7.34 (d, *J* 9.1 Hz, 1H), 7.28 (d, *J* 9.1 Hz, 1H), 7.01 (d, *J* 8.8 Hz, 2H), 3.90 (s, 3H), 2.14 (s, 3H), 2.07 (s, 3H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ 190.0, 182.7, 165.0, 164.1, 159.9, 146.0, 142.9, 142.1, 133.0, 132.6, 125.3, 122.5, 121.9, 114.8, 114.0, 55.6, 13.28, 12.3 ppm; FT-IR: \bar{v}_{max} 2928, 2845, 1788, 1721, 1638, 1610, 1582, 1470, 1263, 1243, 1219, 1058, 1028 cm⁻¹; HRMS: calc for C₂₀H₁₆O₆ [M-H]⁺: 351.0869; found 351.0875.

6,7-Dimethyl-5,8-dioxo-5,8-dihydronaphthalene-1,4-diyl bis(4-nitrobenzoate) The (16). bis-pnitrobenzoylester DMN 16 was synthesized following the general procedure-1 using DMN 3 (50 mg, 0.23 mmol) and p-nitrobenzoyl chloride 11 (0.17 g, 0.92 mmol) in dry THF (5.0 mL) and dry Et₃N (96 µl, 0.69 mmol). The reaction mixture was stirred at 25 °C for 22 hours. The resulting reaction solvent was then removed by rotary evaporator and the crude material was subjected to column chromatography eluting with (EtOAc:Hexane (20:80)) to give the compound **16** (19 mg, 21%) as yellow colour solid along with mono-ester **19** (23 mg, 27%) as orange-coloured solid. An analytical sample of **16** was obtained by recrystallization from dichloromethane/hexane to give orange-colored needles. R_f 0.42 (in dichloromethane); mp 259-261 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.45 (d, J 8.3 Hz, 4H), 8.41 (d, J 8.2 Hz, 4H), 7.56 (s, 2H), 2.04 (s, 6H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ 182.9, 163.4, 151.1, 147.4, 143.8, 134.8, 131.7, 130.4, 124.8, 123.9, 13.0 ppm; FT-IR: \bar{v}_{max} 3113, 1742, 1657, 1526, 1348, 1323, 1267, 1221, 1084, 842, 707 cm⁻¹; HRMS: calc for C₂₆H₁₆N₂O₁₀ [M]⁺: 516.0805; found 516.0811.

4-Hydroxy-6,7-dimethyl-5,8-dioxo-5,8-dihydronaphthalen-1-yl 4-nitrobenzoate (19). R_f 0.53 (in dichloromethane); mp 198-200 °C; ¹H NMR (500 MHz, CDCl₃) δ 12.65 (s, 1H), 8.44 – 8.40 (m, 2H), 8.40 – 8.36 (m, 2H), 7.37 (d, *J* 9.1 Hz, 1H), 7.32 (d, *J* 9.1 Hz, 1H), 2.16 (s, 3H), 2.06 (s, 3H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ 189.8, 182.8, 163.6, 160.2, 151.0, 145.82, 142.6, 142.0, 135.0, 132.3, 131.6, 125.6, 123.8, 122.1, 114.9, 13.2, 12.4 ppm; FT-IR: $\bar{\nu}_{max}$ 3118, 2922, 2853, 1731, 1664, 1638, 1617, 1520, 1470, 1349, 1262, 1243, 1225, 1095, 1064, 850, 709 cm⁻¹; HRMS: calc for C₁₉H₁₂NO₇ [M-H]⁺: 366.0614; found 366.0619.

4-Hydroxy-6,7-dimethyl-5,8-dioxo-5,8-dihydronaphthalen-1-yl acetate (6). DMN **3** (0.10 g, 0.46 mmol) was dissolved in dry THF (10 mL) and one and a half equivalents of dry Et₃N (96 μl, 0.69 mmol) was added slowly. The reaction mixture was stirred for 30 minutes at room temperature and then one equivalent of acetyl chloride **4** (33 μL, 0.46 mmol) was added slowly. The resulting reaction mixture was stirred at 25 °C for 20 hours. The resulting reaction solvent was then removed by rotary evaporator and the crude material was subjected to column chromatography (EtOAc:Hexane (20:80)) to give the target mono-ester compound **6** (43 mg, 36%) as an orange-coloured solid, along with the bis-ester **5** by-product (20 mg, 29%) as yellow-coloured solid. An analytical sample of **6** was obtained by recrystallization from dichloromethane/hexane to give orange color needles. R_f 0.37 (in ethylacetate:hexane (20:80)); mp 143-145 °C; ¹H NMR (500 MHz, CDCl₃) δ 12.62 (s, 1H), 7.24 (s, 1H), 7.23 (s, 1H), 2.42 (s, 3H), 2.15 (s, 3H), 2.12 (s, 3H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ 189.9, 182.9, 169.8, 159.9, 145.8, 142.6, 142.3, 132.6, 125.5, 122.1, 114.8, 21.2, 13.2, 12.3 ppm; FT-IR: \bar{v}_{max} 2920, 2850, 1757, 1652, 1634, 1618, 1461, 1435, 1359, 1280, 1186, 1011, 786 cm⁻¹; HRMS: calc for C₁₄H₁₂O₅ [M]⁺: 260.0685; found 260.0686.

4-Acetoxy-6,7-dimethyl-5,8-dioxo-5,8-dihydronaphthalen-1-yl benzoate (20). To a stirred mixture of monoasetil-DMN **6** (40 mg, 0.15 mmol) in dry THF (3.0 mL) was added one and a half equivalents of dry Et₃N (32 µl, 0.23 mmol) dropwise at 25 °C and stirred for 30 minutes. The reaction mixture was cooled to 0 °C and one and a half equivalents of benzoyl chloride **7** (27 µl, 0.23 mmol) was slowly added. The reaction mixture was warmed to 25 °C and stirred for 2 days. The resulting reaction solvent was then removed by rotary evaporator and the crude material was subjected to column chromatography eluting with dichloromethane to give the compound **20** (9.5 mg, 17%) as yellow-coloured solid. An analytical sample of **20** was obtained by recrystallization from dichloromethane/hexane to give yellow color needles. R_f 0.28 ((in ethylacetate:hexane (20:80)); mp 190-192 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.27 – 8.23 (m, 2H), 7.70 – 7.64 (m, 1H), 7.55 (t, *J* 7.8 Hz, 2H), 7.45 (d, *J* 8.8 Hz, 1H), 7.37 (d, *J* 8.8 Hz, 1H), 2.46 (s, 3H), 2.09 (s, 3H), 2.04 (s, 3H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ 183.3, 183.1, 169.6, 165.1, 147.6, 147.4, 143.6, 143.4, 133.8, 130.6, 130.5, 130.3, 129.4, 128.8, 125.0, 124.8, 21.2, 13.0, 13.0 ppm; FT-IR: \bar{v}_{max} 2919, 1758, 1729, 1658, 1630, 1587, 1449, 1324, 1253, 1220, 1185, 1023, 992, 705 cm⁻¹; HRMS: calc for C₂₁H₁₆O₆ [M+H]⁺: 365.1025; found 365.1014.

Acknowledgements

M. F. S. thanks Sebnem Abadan for her kind assistance in the laboratory, Sadık Metin Ceyhan for helping in analyzing X-ray data, Ibrahim Fazil Sengul for proofreading the manuscript, and Ahmet Ceyhan Gören for his help for HRMS.

Supplementary Material

NMR spectra are provided in the supplementary material file. Additional crystallographic data with CCDC reference numbers 1948087 (5), 1946392 (6), 2048115 (12), 2184617 (19), and 2184607 (20) have been deposited within the Cambridge Crystallographic Data Center via <u>www.ccdc.cam.ac.uk/deposit</u>

References

- 1. Krauth-Siegel, R. L. ; Coombs, G. H. *Parasitol. Today* **1999**, *15*, 404. https://doi.org/10.1016/S0169-4758(99)01516-1
- 2. Segura-Aguilar, J. ; Jönsson, K. ; Tidefelt, U. ; Paul, C. *Leuk. Res.* **1992**, *16*, 631. https://doi.org/10.1016/0145-2126(92)90013-W
- 3. Baik, K-U.; Song, G. Y.; Kim, Y.; Sok, D-E.; Ahn, B-Z. *Arch. Pharm. Med. Chem.* **1997**, *330*, 377. https://doi.org/10.1002/ardp.19973301204
- 4. Valipour, M. *Eur. J. Med. Chem.* **2022**, *235*, 114314. https://doi.org/10.1016/j.ejmech.2022.114314
- Guo, C.; He, J.; Song, X.; Tan, L.; Wang, M.; Jiang, P.; Li, Y.; Cao, Z.; Peng, C. *Pharmacol. Res.* 2019, 149, 104463. https://doi.org/10.1016/j.phrs.2019.104463

Page 210

- 6. Wang, R.; Yin, R.; Zhou, W.; Xu, D.; Li, S *Expert. Opin. Ther. Pat.* **2012**, *22*, 977. <u>https://doi.org/10.1517/13543776.2012.709237</u>
- Sun, W.; Du, L. D.; Wang, S. B.; Chen, X. P.; Du, G. H. In *Natural Small Molecule Drugs from Plants*; Du, G. H. (ed); Springer : Singapore, 2018 ; p 731. <u>https://doi.org/10.1007/978-981-10-8022-7</u>
- 8. Paull, K. D.; Zee-Cheng, R. K. Y.; Cheng, C. C. J. Med. Chem. 1976, 19, 337. https://doi.org/10.1021/jm00224a029
- Rodríguez, J. G.; Smith-Verdier, P.; Florencio, F.; García-Blanco, S. J. Mol. Struct. 1984, 112, 101. <u>https://doi.org/10.1016/0022-2860(84)80247-1</u>
- 10. Dessolin, J.; Biot, C.; Davioud-Charvet, E. *J. Org. Chem.* **2001**, *66*, 5616. <u>https://doi.org/10.1021/jo010137n</u>
- 11. Lu, Q.; Liu, W.; Ding, J.; Cai, J.; Duan, W. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 1375. https://doi.org/10.1016/S0960-894X(02)00158-0
- 12. Wu, X.; Xu, L.; Cai, J. *Bioorg. Med. Chem. Lett.* **1999**, *9*, 2635. https://doi.org/10.1016/S0960-894X(99)00457-6
- 13. Rodriguez, J. G.; Pablo, A. D.; Smith-Verdier, P.; Florencio, F.; Garcia-Blanco, S. *Bull. Chem. Soc. Jpn.* **1986**, *59*, 3957.

https://doi.org/10.1246/bcsj.59.3957

This paper is an open access article distributed under the terms of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/)